

# Wavelengths and oscillator strengths of Xe II from the UVES spectra of four HgMn stars

K. Yüce<sup>1,2</sup>, F. Castelli<sup>1</sup>, and S. Hubrig<sup>3</sup>

<sup>1</sup> Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Trieste, Via Tiepolo 11, I-34143 Trieste, Italy

e-mail: yuce@oats.inaf.it

<sup>2</sup> Ankara University, Faculty of Science, Department of Astronomy and Space Sciences, TR-06100, Tandoğan, Ankara, Turkey

<sup>3</sup> Astrophysical Institute Potsdam, An der Sternwarte 16, D-14482 Potsdam, Germany

## ABSTRACT

**Aims.** In spite of large overabundances of Xe II observed in numerous mercury-manganese (HgMn) stars, Xe II oscillator strengths are only available for a very limited number of transitions. As a consequence, several unidentified lines in the spectra of HgMn stars could be due to Xe II. In addition, some predicted Xe II lines are redshifted by about 0.1 Å from stellar unidentified lines, raising the question about the wavelength accuracy of the Xe II line data available in the literature. For these reasons we investigated the Xe II lines lying in the 3900–4521 Å, 4769–7542 Å, and 7660–8000 Å spectral ranges of four well-studied HgMn stars.

**Methods.** We compared the Xe II wavelengths listed in the NIST database with the position of the lines observed in the high-resolution UVES spectrum of the xenon-overabundant, slowly rotating HgMn star HR 6000, and we modified them when needed. We derived astrophysical oscillator strengths for all the Xe II observed lines and compared them with the literature values, when available. We checked the stellar atomic data derived from HR 6000 by using them to compute synthetic spectra for three other xenon-overabundant, slowly rotating HgMn stars, HD 71066, 46 Aql, and HD 175640. In this framework, we performed a complete abundance analysis of HD 71066, while we relied on our previous works for the other stars.

**Results.** We find that all the lines with wavelengths related to the 6d and 7s energy levels have a corresponding unidentified spectral line, blueshifted by the same quantity of about 0.1 Å in all the four stars, so that we identified these lines as coming from Xe II and modified their NIST wavelength value according to the observed stellar value. We find that the Xe II stellar oscillator strengths may differ from one star to another from 0.0 dex to 0.3 dex. We adopted the average of the oscillator strengths derived from the four stars as final astrophysical oscillator strength.

**Key words.** line:identification-atomic data-stars:atmospheres-stars:chemically peculiar- stars:individual:HR 6000, HD 71066, 46 Aql, HD 175640

## 1. Introduction

Several studies of mercury-manganese (HgMn) stars have pointed out the presence of xenon with overabundances up to 5 dex relative to the solar value  $\log(N_{Xe}/N_{tot}) = -9.87$  (Grevesse & Sauval 1998). This is, for instance, the case of  $\kappa$ Cnc and 33 Gem, for which abundances equal to  $-4.87 \pm 0.13$  dex and  $-4.90 \pm 0.07$  dex were determined by Dworetsky et al. (2008). The xenon overabundance implies the presence of numerous Xe II lines in the spectra of the HgMn stars, but the Xe II transition probabilities are very incomplete, when we compare the large number of transitions listed in the NIST database and the small number of them with an associated  $\log gf$ -value. As a consequence, the computed spectra do not include numerous Xe II lines, raising the doubt that some unidentified lines could just be due to Xe II. In addition, we noticed that the wave-

lengths of several Xe II lines are close, but not coincident with the wavelength of some unidentified stellar lines (Castelli & Hubrig 2007), so that we wondered about the accuracy of the wavelength determination from laboratory spectra.

The most complete work on Xe II is that of Hansen & Persson (1987), who analyzed all the published (Boyce 1936; Humphreys 1939) and unpublished Xe II lines from 392 Å to 10220 Å obtained in laboratory by Humphreys and Boyce. In their discussion on the wavelength accuracy, Hansen & Persson (1987) pointed out that the wavelength accuracy for many lines is too low to be satisfactory, mostly owing to the widely varying quality of the experimental data they used. They announced new experimental work to improve the Xe II atomic data. Unfortunately, this work has never been published up to now, all the more so that some preliminary results had indicated that, for the high 6d and 7s levels, there were shifts of about  $0.5 \text{ cm}^{-1}$  between the energy levels determined from the

Humphrey wavelengths and the energy levels determined from the new data. This energy difference corresponds to a difference of 0.1 Å in wavelengths.

Saloman (2004), who performed a critical compilation of all the work on energy levels and wavelengths of Xe II made up to that time, adopted the data from Hansen & Persson (1987) for almost all the lines of the optical region. The Saloman (2004) critical compilation is the one adopted by the NIST database.

To study wavelengths and log *gf*-values of the Xe II lines having intensities  $\geq 100$  in the NIST line list, we used UVES spectra of the four xenon overabundant HgMn stars HR 6000, HD 71066, 46 Aql, and HD 175640. They are slowly rotating stars with  $v\sin i$  1.5 km s $^{-1}$ , 1.5 km s $^{-1}$ , 1.0 km s $^{-1}$ , and 2.5 km s $^{-1}$ , respectively.

We already performed a complete abundance analysis for HD 175640 (Castelli & Hubrig 2004a)<sup>1</sup> and for HR 6000 and 46 Aql (Castelli et al. 2009)<sup>2</sup>. To be consistent with the other papers, we present here an abundance analysis of HD 71066, which was studied with the same methods as adopted for the other stars. A previous work on HD 71066, related to vertical abundance stratification in HgMn stars, was performed by Thiam et al. (2010), who adopted the same observations as are used in this paper. We note, however, that no mention about Xe II was made in their study.

## 2. Observations and data reduction

All the stars were observed at the European Southern Observatory (ESO) using the Very Large Telescope Ultraviolet and Visible Echelle Spectrograph (UVES) with a resolving power ranging from 80000 to 110000.

HD 175640 was observed on June 13, 2001 (Castelli & Hubrig 2004a). HR 6000, 46 Aql, and HD 71066 were part of the same observational run (ESO program 076.D-0169(A)). The spectra of HR 6000 were observed on September 19, 2005, those of 46 Aql on October 18, 2005 (Castelli et al., 2009), while the spectra of HD 71066 were taken on October 27, 2005. Because Nunez et al. (2010) found spectral variations in 19 HgMn stars out of a sample of 28 HgMn stars analyzed, we investigate about a possible variability of HD 71066 by comparing the spectrum observed in 2005 with an UVES spectrum observed in April 2004. We did not find any clear indication of variability.

The spectra of the four stars cover the region 3030 – 10000 Å. For HD 175640 there are two gaps at  $\lambda\lambda$  5759 – 5835 Å and 8519 – 8656 Å. For the other three stars, the gaps occur at 4520 – 4769 Å and 7536 – 7660 Å. All the spectra were reduced by the UVES pipeline Data Reduction Software (Ballester et al. 2000). We analyzed flux-calibrated spectra for the 3050–5750 Å region and RED\_SCI\_POINT spectra for the 5750–9460 Å interval, in that flux-calibrated reduction for the red spectra was not implemented in the pipeline reduction procedure.

The measurement procedures on the spectra of HD 175640, HR 6000, and 46 Aql were described in Castelli & Hubrig (2004a) and Castelli & Hubrig (2007). The spectra of HD 71066 were normalized to the continuum using the IRAF continuum task. The equivalent widths were measured by a Gaussian fitting using the IRAF splot task.

The signal-to-noise ratio (S/N) is different for the different stars. In the spectra of HD 175640, it ranges from 200 in the near UV to 400 in the visual region. It is higher than the S/N of the spectra of the other three stars, which were observed in a different epoch. Furthermore, for each star, it is different in the different spectral intervals. For instance, for HR 6000, it is about 100 in the 5800–6800 Å interval and lowers to about 25 at 7400 Å (REDL spectrum). It is about 50 at 7800 Å and decreases to about 25 at 9400 Å (REDU spectrum). This behavior is similar for 46 Aql and HD 71066. At 6800 Å the S/N is about 100 for HR 6000, 70 for 46 Aql, 100 for HD 71066, and 125 for HD 175640.

## 3. The HgMn star HD 71066

Previous studies of HD 71066 ( $\kappa^2$  Vol, HR 3302) have pointed out the isotopic anomaly of Hg (Dolk et al. 2003; Thiam et al. 2010). No vertical abundance stratification for Ti, Cr, and Fe is found by Thiam et al. (2010). No presence of magnetic field is found both from the inspection of the equivalent widths of the Fe II lines at 6147.7 Å and 6149.2 Å (Hubrig et al. 1999) and after using the FORS 1 spectropolarimeter at the VLT (Hubrig et al. 2006).

### 3.1. Model parameters and abundances of HD 71066

The starting model parameters of HD 71066,  $T_{\text{eff}}=12045$  K, and  $\log g=3.9$  were derived both from the Strömgren photometry and the Fe I – Fe II ionization equilibrium constraint.

The observed colors  $(b-y)=-0.053$ ,  $m=0.122$ ,  $c=0.731$ ,  $\beta=2.769$  were taken from the Hauck & Mermilliod (1998) Catalogue<sup>3</sup>. The synthetic colors were taken from the grid computed by Castelli for  $[M/H]=0$  and microturbulent velocity  $\xi=0$  km sec $^{-1}$ <sup>4</sup>. Zero reddening was adopted for this star, in agreement with the results from the UVBYLIST code of Moon (1985). Observed  $c$  and  $\beta$  indices are reproduced by synthetic indices for model parameters  $T_{\text{eff}}=12045$  K and  $\log g=3.9$

The parameters from the photometry were used for computing an ATLAS9 model with solar abundances for all the elements and zero microturbulent velocity. Using the WIDTH code (Kurucz 2005), we derived the Fe I and Fe II abundance from the equivalent widths of 12 Fe I lines and 26 Fe II lines. Seven of the Fe II lines are transitions between high-excitation energy levels, and they have experimental log *gf*-values. They were used to determine the iron abundance, in that they are rather independent of  $T_{\text{eff}}$  and  $\log g$  (Castelli et al., 2009). Then, we searched for the model atmosphere giving this same abundance from both Fe I lines and low-excitation Fe II lines. All the adopted lines are listed in Table A.1 of Appendix A (on-

<sup>1</sup> <http://wwwuser.oat.ts.astro.it/castelli/hd175640/hd175640.html>

<sup>2</sup> <http://wwwuser.oat.ts.astro.it/castelli/hr6000new/hr6000.html>


<sup>3</sup> <http://obswww.unige.ch/gcpd/gcpd.html>

<sup>4</sup> <http://wwwuser.oat.ts.astro.it/castelli/colors/uvbybeta.html>









**Fig. 4.** Comparison of astrophysical log  $gf$ -values of Xe II derived from HR 6000 with the astrophysical log  $gf$ -values derived from HD 175640.

in Table 4. Wavelengths and log  $gf$ -values were taken from the NIST database (Version 4)<sup>6</sup>. We assumed the classical radiative damping constant  $\gamma_R = 0.2223 \times 10^{16}/\lambda^2$  s $^{-1}$  for  $\lambda$  in Å. For Stark broadening we used the experimental results from Djurovic et al. (2006). Because they are given for a temperature of T=22000 K, we investigated the effect of the temperature on the Stark damping constant  $\gamma_S$ . We interpolated for T=12000 K in the tables from Popovic & Dimitrijevic (1996), which list Stark widths computed at different temperatures. The last two columns of Table 4 compare  $\gamma_S$  values from Djurovic et al. (2006) (Dj) with the interpolated values for temperature from Popovic & Dimitrijevic (1996) (PD). We found that the differences in  $\gamma_S$  from the two sources do not affect the abundances more than 0.01 dex. The approximations of the WIDTH code were used (Castelli 2005) for no available Stark damping constants and for Van der Waals damping constants.

The measured equivalent widths of the selected Xe II lines and the corresponding abundances are given in Table 5.

## 5. Stellar wavelengths and the astrophysical log $gf$ -values for Xe II

Because xenon is more abundant in HR 6000 than in the other stars, we searched in the HR 6000 spectra for those Xe II lines with an intensity equal to or higher than 100 in the NIST line list. When these lines were observed in the spectra, they were added in our line list. For lines with no available log  $gf$ -values, we assigned guessed values based on the line intensity. We examined the interval 3900–8000 Å with two gaps in the 4525–4780 Å and 7536 – 7660 Å regions, due to the lack of spectra in these ranges.

A synthetic spectrum for HR 6000 was computed for the xenon abundance given in Table 5 and for the abundances of all the other elements as derived by Castelli et al. (2009). In all the stars, the wavelength scale was fixed by shifting the observed spectrum on the computed spectrum until overimposing some lines with well-determined wavelength values such as

Ca II 3933.663 Å, Mg II 4481.126 Å, 4481.150 Å, 4481.325 Å, and several strong Fe II lines.

For all the considered Xe II lines, we adjusted the log  $gf$ -value until the observed and computed profiles agree best. For several lines we also adjusted the NIST wavelength, because we noticed that, while they do not have an observed counterpart, they are close to an unidentified stellar line with wavelength blueshifted up to 0.1 Å from the predicted Xe II line.


The astrophysical log  $gf$ -values and the adjusted wavelengths were then checked on the three other stars by comparing their observed spectra with synthetic spectra computed with the Xe II wavelengths and oscillator strengths obtained from the spectrum of HR 6000. The Xe II abundances adopted for the three stars are those given in Table 5. Table B.1 in Appendix B lists wavelengths and log  $gf$ -values as derived from the four stars. We found that for all the examined transitions, the stellar wavelength is the same in the four stars, except for the lines at 5260.44 Å and 6343.96 Å. The largest difference between stellar and NIST wavelength is −0.13 Å observed for the line at 4330.52 Å. This line, as well as all the other lines with  $\Delta\lambda \sim 0.1$  Å has a 6d or a 7s level as upper level. The uncertainty of the energy of these levels is on the order of 0.5 cm $^{-1}$  according to Hansen & Persson (1987).

Figures 1, 2, 3, and 4 show the comparison of the HR 6000 astrophysical log  $gf$ -values for Xe II with the log  $gf$ -values taken from the NIST database and with the astrophysical log  $gf$ -values derived from the spectra of HD 71066, 46 Aql, and HD 175640, respectively (Table B.1, col. 5). The largest discrepancy with the NIST data occurs for the line at 4414.84 Å. We adopted the stellar log  $gf$ -value for it because it gives an excellent agreement between the observed and computed profiles in all the four stars we examined. The comparison of the astrophysical log  $gf$ -values of HR 6000 with those from the other stars shows that they are on average lower by about 0.04–0.05 dex than those from the other stars and that the mean square deviation from the average increases with the decrease in the xenon abundance. We note that the weaker a line, the more uncertain its astrophysical log  $gf$ -value, mostly when the noise is not negligible. In particular, red spectra are affected both by rather large noise and by numerous telluric lines that lower the accuracy of the results.

The final line list for Xe II is shown in Table 6. Columns 1 and 2 give the wavelength derived from the stellar spectra and the astrophysical log  $gf$ -value obtained by averaging the astrophysical log  $gf$ -values from the four stars. The associated error is the standard deviation from the mean. When it is not given, it means that the log  $gf$ -value was obtained from only one star. Columns 3 and 4 list log  $gf$ -values from the literature and the source. The literature sources are the NIST database, version 4 (NIST4), and Zieleńska et al. (2002) (ZBD). Zieleńska et al. (2002) estimate that, in general, their experimental transition rates agree with the NIST critical compilation made by Reader et al. (1980), which is the one adopted in the NIST4 database.

The last column gives the  $\gamma_{Stark}$  parameter, which was determined as described in Sect. 4. Figure 5 shows, for each studied star, the comparison of the observed and computed spectra in the region of the Xe II line with wavelength 4462.190 Å, according to the NIST database, and 4462.090 Å, according to

<sup>6</sup> <http://www.nist.gov/pml/data/asd.cfm>



**Fig. 5.** Comparison for the four stars of the observed (black line) and computed (red line) spectra in the region of the Xe II line with wavelength 446.2190 nm according to the NIST database (left panel) and with wavelength 446.2090 nm according to this paper (right panel). The line identification can be decoded as follows: for the first line, 898 last 3 digits of wavelength 446.0898 nm; 26 atomic number of iron; .01 charge/100; i.e., 26.01 identifies the line as Fe II; 91 070 is the energy of the lower level in  $\text{cm}^{-1}$ ; 919 is the residual central intensity in per mil. The Xe II line is identified by 54.01.

Table 6. The wavelength shift of  $0.1 \text{ \AA}$  and the astrophysical  $\log g f$  value of  $+0.33$ , which are the same for all the stars, provide excellent agreement between the observed and computed Xe II profiles.

## 6. Conclusions

From the high resolution stellar spectra of four HgMn stars we derived both wavelengths and  $\log g f$ -values for 100 Xe II lines, which should also be observable in the spectra of numerous others chemically peculiar B-type stars. Of these lines, only 22 lines have  $\log g f$ -values available in the NIST database. The NIST wavelength of two of them,  $4180.10 \text{ \AA}$  and  $4330.52 \text{ \AA}$ ,

differs by about  $0.1 \text{ \AA}$  from that observed in the spectra. There is a total of 27 lines in our sample for which the observed wavelength differs from the NIST wavelength by more than  $-0.06 \text{ \AA}$  with the maximum shift of  $-0.13 \text{ \AA}$  for the line at  $4330.52 \text{ \AA}$ . We believe that the wavelength differences are mostly the result of uncorrect energy levels, in that they are all related to 6d or 7s levels, which have an uncertainty of about  $0.5 \text{ cm}^{-1}$  according to Hansen & Persson (1987). This hypothesis seems us to be more reliable than that of some isotopic anomaly for Xe. For instance, using the isotopic wavelengths from Alvarez et al. (1979), Castelli & Hubrig (2007) excluded that the blueshift of  $0.03 \text{ \AA}$  observed for the Xe II line at  $6051.15 \text{ \AA}$  can be due to some isotopic anomaly. Instead, because no isotopic compo-





**Table 7.** A few 7s, 5d, and 6d even Xe II energy levels from Hansen & Persson (1987) modified according to the wavelength positions observed in the UVES spectra of HR 6000, HD 71066, 46 Aql, and HD 175640

Term	level value (cm <sup>-1</sup> )		
	NIST	This paper	
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>3</sup> P <sub>2</sub> )7s	[2] <sub>5/2</sub>	132518.82	132519.23
	[2] <sub>3/2</sub>	133189.42	133189.94
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>3</sup> P <sub>0</sub> )7s	[0] <sub>1/2</sub>	140883.42	140883.79
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>1</sup> D <sub>2</sub> )5d	[0] <sub>1/2</sub>	135060.97	135061.36
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>3</sup> P <sub>2</sub> )6d	[4] <sub>9/2</sub>	136109.65	136110.13
	[4] <sub>7/2</sub>	136597.81	136598.48
	[3] <sub>7/2</sub>	135507.32	135507.72
	[3] <sub>5/2</sub>	139094.28	139094.83
	[2] <sub>5/2</sub>	135547.13	135547.53
	[2] <sub>3/2</sub>	135708.32	135708.72
	[1] <sub>3/2</sub>	139640.43	139640.61
	[1] <sub>1/2</sub>	136554.11	136554.47
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>3</sup> P <sub>1</sub> )6d	[3] <sub>7/2</sub>	145587.61	145588.12
	[3] <sub>5/2</sub>	146927.86	146928.34
	[2] <sub>3/2</sub>	145940.34	145940.79
	[1] <sub>3/2</sub>	148085.19	148085.36
	[1] <sub>1/2</sub>	145222.72	145223.16
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>3</sup> P <sub>0</sub> )6d	[2] <sub>5/2</sub>	144384.90	144385.45
	[2] <sub>3/2</sub>	144140.16	144140.69
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>1</sup> D <sub>2</sub> )6d	[4] <sub>9/2</sub>	152806.73	152806.73 ?
	[4] <sub>7/2</sub>	152708.92	152709.19
	[1] <sub>3/2</sub>	153584.09	153584.02

# **Online Material**

## **Appendix A: The lines used for the abundance analysis of HD 71066**

Table A.1 lists the lines that were used to derive the abundances of HD 71066. The wording “not obs” is given for lines not present in the spectra, while the wordings “profile” and “blend” are given for lines observed well in the spectra, but that do not have measurable equivalent widths either because the noise affects the profile too much or because other components affect the line. These wordings also indicate lines for which adequate equivalent widths cannot be computed, as in the cases of Mg II at 4481 Å which is a blend of transitions belonging to the same multiplet, of most Mn II lines that are affected by hyperfine structure, of the Ca II infrared triplet, which is a blend of isotopic components, and so on. For the remaining lines the measured equivalent widths are given in the table.

## **Appendix B: The investigated Xe II lines in HR 6000, HD 71066, 46 Aql, and HD 175640**

Table B.1 gives the details on the determination of the Xe II wavelengths and log *gf*-values from the spectra of the four stars. It lists in successive columns the laboratory wavelengths and the line intensity taken from the NIST database (footnote 6), the stellar wavelengths as derived from HR 6000, HD 71066, 46 Aql, and HD 175640. If the observed wavelength is the same in all the stars, only that of HR 6000 is given. HR 6000, HD 71066, 46 Aql, and HD 175640 are indicated in col. 6 with the numbers 1,2, 3, and 4, respectively. A question mark means uncertain determinations from that star. The wavelength difference  $\Delta\lambda = \lambda(\text{stellar}) - \lambda(\text{lab})$  is given in col. 4. The energy and the configuration of the lower and upper level of the transition are given in cols. 7,8,9, and 10, respectively. The last column adds some notes about the observed lines. Table B.1 lists also the NIST log *gf*-values and the log *gf*-values derived from the experimental transition rates determined by Zieleńska et al. (2002).

**Table A.1.** Abundances of HD 71066 from the ATLAS12 model with parameters  $T_{\text{eff}}=12000$  K,  $\log g=4.1$ 

HD 71066[12000,4.1,AT12]							
Species	$\lambda(\text{\AA})$	$\log gf$	Ref. <sup>a</sup>	$\chi_{\text{low}}$	W(m\AA)	$\log(N_Z)/N_{\text{tot}}$	Notes
He I <sup>a</sup>	4026.209	-0.374	NIST4	169087.008	profile	-2.28	the core is computed too strong
He I	4471.502	+0.043	NIST4	169087.008	profile	-2.28	the core is computed too strong
He I	5875.661	+0.739	NIST4	169086.964	profile	-2.50	the core is computed too strong
He I	6678.151	+0.328	NIST4	171135.00	profile	-2.50	
Be II	3130.420	-0.178	NIST4	0.00	profile	-10.80	
C II	3918.968	-0.533	NIST4	131724.370	profile	-3.90	observed at 3918.92
C II	4267.001	+0.563	NIST4	145549.270	profile	-3.90	observed at 4267.10
C II	4267.261	+0.716	NIST4	145550.700	profile	-3.90	
C II	4267.261	-0.584	NIST4	145550.700	profile	-3.90	
C II	6578.052	-0.021	NIST4	116537.65	profile	-3.90	
C II	7236.420	+0.294	NIST4	131735.52	profile	-3.90	observed at 7236.35
N I	8680.282	+0.359	NIST4	83364.620	not obs	$\leq -5.50$	
N I	8683.403	+0.105	NIST4	88317.830	not obs	$\leq -5.50$	
O I	4368.193	-2.665	NIST4	76794.978	profile	-3.70	
O I	4368.242	-1.964	NIST4	76794.978	profile	-3.70	
O I	4368.258	-2.186	NIST4	76794.978	profile	-3.70	
O I	5329.096	-1.938	NIST4	86625.757	profile	-3.67:	
O I	5329.099	-1.586	NIST4	86625.757	profile	-3.67:	
O I	5329.107	-1.695	NIST4	86625.757	profile	-3.67:	
O I	6155.961	-1.363	NIST4	86625.757	profile	-3.57	
O I	6155.971	-1.011	NIST4	86625.757	profile	-3.57	
O I	6155.989	-1.120	NIST4	86625.757	profile	-3.57	
O I	6156.737	-1.487	NIST4	86627.778	profile	-3.57	
O I	6156.755	-0.898	NIST4	86627.778	profile	-3.57	
O I	6156.778	-0.694	NIST4	86627.778	profile	-3.57	
O I	6454.444	-1.066	NIST4	86627.778	profile	-3.57	
O I	6455.977	-0.920	NIST4	86631.454	profile	-3.60	
O I	7002.173	-2.644	NIST4	88631.146	profile	-3.58	
O I	7002.196	-1.489	NIST4	88631.146	profile	-3.58	
O I	7002.230	-0.741	NIST4	88631.146	profile	-3.58	
O I	7002.250	-1.364	NIST4	88631.303	profile	-3.58	
Ne I	6402.248	+0.345	NIST4	134041.840	not obs	$\leq -5.70$	
Ne I	7032.413	-0.249	NIST4	134041.840	not obs	$\leq -5.70$	
Na I	5889.950	+0.108	NIST4	0.00	38.2	-5.42	
Na I	5895.924	-0.194	NIST4	0.00	19.3	-5.62	

**Table A.1.** cont.

Species	$\lambda(\text{\AA})$	$\log gf$	Ref. <sup>a</sup>	HD 71066[12000,4.1,AT12]			
				$\chi_{low}$	W(m\AA)	$\log(N_Z)/N_{tot}$	Notes
Mg I	5167.321	-0.870	NIST4	21850.405	1.30	-5.36	
Mg I	5172.684	-0.393	NIST4	21870.464	4.60	-5.27	
Mg II	4481.126	+0.749	NIST4	71490.190	profile	-5.40	
Mg II	4481.150	-0.553	NIST4	71490.190	profile	-5.40	
Mg II	4481.325	+0.594	NIST4	71491.063	profile	-5.40	
Al I	3944.006	-0.638	NIST4	0.000	not obs	$\leq -7.30$	
Al I	3961.520	-0.336	NIST4	112.061	not obs	$\leq -7.30$	
Al II	7056.712	+0.110	NIST4	91274.500	not obs	$\leq -7.30$	
Si II	3853.665	-1.341	NIST4	55309.350	66.7	-4.81	
Si II	3856.018	-0.406	NIST4	55325.180	113.7	-4.91	
Si II	3862.595	-0.757	NIST4	55309.350	101.4	-4.74	
Si II	4072.709	-2.701	NIST4	79338.500	2.3	-4.32	
Si II	4075.452	-1.400	NIST4	79355.020	16.37	-4.65	
Si II	4190.724	-0.351	LA	108820.600	8.35	-4.58	
Si II	4198.133	-0.611	LA	108778.700	5.98	-4.48	
Si II	5041.024	+0.029	NIST4	81191.340	82.89	-4.35	
Si II	5055.984	+0.523	NIST4	81251.320	100.6	-4.60	
Si II	5056.317	-0.492	NIST4	81251.320	46.08	-4.48	
Si II	5957.559	-0.225	NIST4	81191.340	44.44	-4.57	
Si II	5978.930	+0.084	NIST4	81251.320	56.53	-4.62	
Si II	7849.722	+0.470	NIST4	101024.350	10.65	-4.94	
P II	4044.576	+0.481	K,MRB	107360.250	18.35	-5.04	
P II	4127.559	-0.110	K,KP	103667.860	8.01	-5.13	
P II	4288.606	-0.630	K,MRB	101635.690	2.10	-5.34	
P II	4420.712	-0.329	NIST4	88893.220	15.97	-5.12	
P II	4452.472	-0.194	K,MRB	105302.170	6.30	-5.00	
P II	4463.027	+0.026	K,MRB	105549.670	8.67	-5.04	
P II	4466.140	-0.560	NIST4	105549.670	1.83	-5.24	
P II	4475.270	+0.440	NIST4	105549.670	13.15	-5.20	
P II	5296.077	-0.160	NIST4	87124.600	22.90	-4.86	
P II	5344.729	-0.390	NIST4	86597.550	15.49	-4.99	
P II	5425.880	+0.180	NIST4	87124.600	31.31	-4.92	
P II	6034.039	-0.220	NIST4	86597.550	16.81	-4.95	
P II	6043.084	+0.416	NIST4	87124.600	32.87	-4.94	

**Table A.1.** cont.

HD 71066[12000,4.1,AT12]						
Species	$\lambda(\text{\AA})$	$\log gf$	Ref. <sup>a</sup>	$\chi_{low}$	W(m\AA)	$\log(N_Z)/N_{tot}$
P III	4222.198	+0.210	NIST4	117835.950	4.99	-5.13
S II	4153.068	+0.617	NIST4	128233.200	2.98	-5.66
S II	4162.665	+0.777	NIST4	128599.160	2.67	-5.87
Ca I	4226.728	+0.244	NIST4	0.000	profile	-5.68
Ca II	3158.869	+0.27	NIST4	25191.51	29.96	-6.60
Ca II	3179.331	+0.52	NIST4	25414.40	32.47	-6.73
Ca II	3181.275	-0.45	NIST4	25414.40	15.90	-6.45
Ca II	3933.663	+0.135	NIST4	0.000	profile	-6.33
Ca II	3968.469	-0.18	NIST4	0.000	profile	-6.90
Ca II	8498.023	-1.45	GAL	13650.19	profile	-6.33
Ca II	8542.091	-0.50	GAL	13710.88	profile	-6.33
Ca II	8662.142	-0.76	GAL	13650.19	profile	-6.33
Sc II	4246.822	+0.242	NIST4	2540.950	not obs	$\leq -10.5$
Sc II	4314.083	-0.100	NIST4	4987.790	not obs	$\leq -10.5$
Ti II	4163.644	-0.130	PTP	20891.660	40.17	-6.45
Ti II	4287.873	-1.790	PTP	8710.440	9.09	-6.51
Ti II	4290.215	-0.850	PTP	9395.710	37.88	-6.51
Ti II	4294.094	-0.930	PTP	9744.250	40.01	-6.41
Ti II	4300.042	-0.440	PTP	9518.060	57.29	-6.39
Ti II	4301.922	-1.150	PTP	9363.620	24.83	-6.55
Ti II	4367.652	-0.860	PTP	20891.660	12.52	-6.53
Ti II	4395.031	-0.540	PTP	8744.250	55.79	-6.38
Ti II	4399.765	-1.190	PTP	9975.920	24.64	-6.94
Ti II	4411.072	-0.670	PTP	24961.030	13.25	-6.48
Ti II	4417.714	-1.190	PTP	9395.710	24.69	-6.44
Ti II	4443.810	-0.720	PTP	8710.440	49.85	-6.37
Ti II	4464.448	-1.810	PTP	9363.620	9.85	-6.42
Ti II	4468.492	-0.620	NIST4	9118.260	51.86	-6.41
Ti II	4488.325	-0.510	PTP	25192.710	16.92	-6.46
Ti II	4805.085	-1.120	NIST4	16625.110	18.79	-6.31
Ti II	4911.195	-0.610	PTP	25192.790	14.40	-6.44
V II	3093.105	+0.559	K10V	3162.800	not obs	$\leq -10.0$
V II	3102.294	+0.434	K10V	2968.220	not obs	$\leq -10.0$

**Table A.1.** cont.

Species	$\lambda(\text{\AA})$	$\log gf$	Ref. <sup>a</sup>	HD 71066[12000,4.1,AT12]		
				$\chi_{low}$	W(m\AA)	$\log(N_Z)/N_{tot}$
Cr II	4812.337	-1.997	K10Cr	31168.580	6.07	-6.22
Cr II	4824.127	-0.980	K10Cr	31219.350	39.36	-6.06
Cr II	4836.229	-1.963	K10Cr	31117.390	7.39	-6.16
Cr II	5237.329	-1.160	NIST4	32854.310	22.48	-6.24
Cr II	5246.768	-2.460	NIST4	29951.880	2.93	-6.17
Mn II	3917.318	-1.135	K09Mn	55759.270	profile	-5.93
Mn II	4363.255 <sup>b</sup>	-1.887	K09Mn	44899.820	profile	-5.93
Mn II	4365.217 <sup>b</sup>	-1.344	K09Mn	44899.820	profile	-5.93
Mn II	4478.637 <sup>b</sup>	-0.945	K09Mn	53597.130	profile	-5.93
Mn II	4806.823	-1.571	K09Mn	43696.120	profile	-6.03
Fe I	3581.193	+0.406	FW06	6928.27	28.28	-3.68
Fe I	3618.768	-0.003	FW06	7985.78	15.49	-3.88
Fe I	4005.242	-0.610	FW06	12560.93	14.40	-3.87
Fe I	4071.738	-0.022	FW06	12698.55	31.00	-3.91
Fe I	4202.029	-0.708	FW06	11976.24	13.44	-3.85
Fe I	4219.360	+0.000	FW06	28819.95	7.80	-3.82
Fe I	4235.936	-0.341	FW06	19562.44	12.27	-3.81
Fe I	4271.760	-0.164	FW06	11976.24	30.68	-3.84
Fe I	4383.545	+0.200	FW06	11976.24	43.10	-3.85
Fe I	4404.750	-0.142	FW06	12560.93	29.22	-3.87
Fe I	4415.122	-0.615	FW06	12968.55	14.49	-3.84
Fe I	5364.871	+0.228	FW06	35856.40	3.77	-3.96
Fe II	4128.748	-3.580	FW06	20830.58	31.85	-3.92
Fe II	4178.862	-2.440	FW06	20830.58	67.36	-3.96
Fe II	4273.326	-3.300	FW06	21812.05	41.90	-3.86
Fe II	4296.572	-2.930	FW06	21812.05	54.16	-3.85
Fe II	4369.411	-3.580	FW06	22409.85	27.75	-3.93
Fe II	4413.601	-4.190	FW06	21581.64	15.83	-3.75
Fe II	4416.830	-2.600	FW06	22409.85	64.47	-3.83
Fe II	4491.405	-2.640	FW06	23031.30	57.57	-3.96
Fe II	4508.288	-2.350	FW06	23031.30	73.82	-3.76
Fe II	4515.339	-2.360	FW06	23939.36	65.01	-3.94
Fe II	4913.295	+0.016	J07	82978.71	33.22	-3.77
Fe II	4993.358	-3.680	FW06	22637.20	26.98	-3.83
Fe II	5001.953	+0.933	J07	82853.65	65.38	-3.85
Fe II	5030.631	+0.431	FW06	82978.68	44.23	-3.87
Fe II	5035.700	+0.630	FW06	82978.68	52.34	-3.84

**Table A.1.** cont.

HD 71066[12000,4.1,AT12]							
Species	$\lambda(\text{\AA})$	$\log g f$	Ref. <sup>a</sup>	$\chi_{low}$	W(m\AA)	$\log(N_Z)/N_{tot}$	Notes
Fe II	5144.352	+0.307	FW06	84424.37	23.98	-4.24	
Fe II	5247.956	+0.550	FW06	84938.18	41.29	-3.88	
Fe II	5260.254	+1.090	J07	84863.38	65.44	-3.84	
Fe II	5276.002	-1.900	FW06	25805.33	76.52	-3.95	
Fe II	5339.592	+0.568	J07	84296.87	44.50	-3.85	
Fe II	5414.852	-0.258	J07	84863.38	20.80	-3.72	
Fe II	5425.257	-3.390	FW06	25805.33	36.19	-3.64	
Fe II	5465.932	+0.348	FW06	85679.70	38.16	-3.70	
Fe II	5493.830	+0.259	FW06	84685.20	33.73	-3.80	
Fe II	5506.199	+0.923	J07	84863.38	53.95	-3.89	
Fe II	5510.783	+0.043	J07	85184.77	27.35	-3.76	
Co II	4160.657	-1.751	K06Co	27484.371	blend	$\leq -7.88$	
Ni II	4067.031	-1.834	K03Ni	32499.530	blend	$\leq -7.90$	
Cu II	4909.734	+0.790	K03Cu	115568.985	not obs	$\leq -7.8$	
Zn II	4911.625	+0.540	NIST4	96909.740	not obs	$\leq -7.94$	
As II	4466.348						
As II	4494.230						
As II	5105.58			81508.925	3.74		
As II	5231.38			79128.330	3.16		
As II	5331.23			81508.925	7.07		
As II	5497.727			78730.893	4.52	blend	
As II	5558.09			79128.330	7.11	blend	
As II	5651.32			81508.925	9.29		
As II	6110.07			82819.214	2.32		
As II	6170.27			79128.330	2.62	blend	
Sr II	4077.709	+0.151	NIST4	0.000	32.45	-8.27	
Y II	3950.349	-0.485	NIST4	840.213	17.38	-7.68	
Y II	4883.682	+0.070	NIST4	8743.316	25.22	-7.49	
Y II	4900.120	-0.090	NIST4	8328.041	20.57	-7.52	

**Table A.1.** cont.

HD 71066[12000,4.1,AT12]							
Species	$\lambda(\text{\AA})$	$\log gf$	Ref. <sup>a</sup>	$\chi_{low}$	W(m\AA)	$\log(N_Z)/N_{tot}$	Notes
Xe II	4844.33	+0.49	NIST4	93068.440	20.72	-5.43	
Xe II	5292.21	+0.35	NIST4	93068.440	19.72	-5.20	
Xe II	5419.14	+0.21	NIST4	95064.38	14.27	-5.24	
Xe II	5438.97	-0.19	NIST4	102799.07	2.93	-5.55	
Xe II	5472.61	-0.45	NIST4	95437.67	5.03	-5.34	
Xe II	5531.06	-0.62	NIST4	95437.67	1.87	-5.71	
Xe II	5719.61	-0.74	NIST4	96033.48	1.40	-5.64	
Xe II	5976.46	-0.22	NIST4	95064.38	4.70	-5.49	
Xe II	6036.20	-0.61	NIST4	95396.74	2.44	-5.45	
Xe II	6051.15	-0.25	NIST4	95437.67	4.59	-5.44	
Xe II	6097.59	-0.24	NIST4	95436.74	3.93	-5.53	
Xe II	6990.88	+0.20	NIST4	99409.99	5.18	-5.36	
Nd III	4927.488	-0.83	DREAM	3715.	1.78	-9.63	
Nd III	5294.113	-0.65	DREAM	0.	4.12	-9.62	
Dy III	3930.640	-0.88	DREAM	0.	profile	-9.90	
Au II	4016.067	-1.88	RW	84510.894	2.39	-7.15	
Au II	4052.790	-1.69	RW	84510.894	3.99	-7.08	
Hg I	4358.314	-0.321	NIST4	39412.300	profile	-6.40	
Hg II	3983.890	-1.51	NIST4	35514.000	profile	-6.40	
Hg II	5677.102	+0.82	NIST4	105543.000	5.56	-6.19	blend

<sup>a</sup> He I profiles were compute as described in Castelli & Hubrig (2004a). The wavelengths and  $\log gf$ -values are multiplet values.

<sup>b</sup> The hyperfine structure was considered in the line profile computations.

DREAM: Biémont et al.(1999): <http://w3.umons.ac.be/astro/dream.shtml>;

NIST4: NIST Atomic Spectra Database, version 4 at <http://physics.nist.gov/pml/data/asd.cfm>;

FW06: Fuhr & Wiese (2006); GAL: Gallagher (1967); LA: Lanz & Artru (1985); PTP: Pickering et al. (2002);

J07: Johansson (2007);

K03Ni: <http://kurucz.harvard.edu/atoms/2801/gf2801.pos>;

K03Cu: <http://kurucz.harvard.edu/atoms/2901/gf2901.pos>;

K06Co: <http://kurucz.harvard.edu/atoms/2701/gf2701.pos>;

K09Mn: <http://kurucz.harvard.edu/atoms/2501/gf2501.pos>;

K10V: <http://kurucz.harvard.edu/atoms/2301/gf2301.pos>;

K10Cr: <http://kurucz.harvard.edu/atoms/2401/gf2401.pos>;

"K" before another  $\log gf$  source means that the  $\log gf$  is from the Kurucz files available at <http://kurucz.harvard.edu/linelists/gf100/>; in particular: KP: Kurucz & Peytremann (1975); MRB: Miller et al. (1971);

RW: Rosberg & Wyart (1997).



**Table B.1.** cont.

$\lambda$ (Lab)	Int.	$\lambda$ (stellar)	$\Delta\lambda$	$\log g f$	$\chi_{low}(\text{cm}^{-1})$	Term	$\chi_{up}(\text{cm}^{-1})$	Term	Notes		
4414.84	300	4414.84	0.00	-0.50 +0.243N	1,2,3	$(^1\text{D}_2)6\text{p}$	[3]7/2	132207.76	$(^1\text{D}_2)6\text{p}$	[2]5/2	2,4 blend
4416.07	150	4416.090	+0.02	-0.80	1 ?	$(^3\text{P}_1)6\text{p}$	[1]3/2	146927.86	$(^3\text{P}_1)6\text{d}$	[3]5/2	3 noise
4448.13	500	4448.025	-0.105	+0.05 +0.15	1,4 2,3	$(^3\text{P}_1)6\text{p}$	[2]5/2	145587.61	$(^3\text{P}_1)6\text{d}$	[3]7/2	
4462.19	1000	4462.090	-0.10	+0.33	1,2,3	$(^3\text{P}_2)6\text{p}$	[3]7/2	136109.65	$(^3\text{P}_2)6\text{d}$	[4]9/2	4 blend
4787.77	100	4787.77	0.00	-0.88 -0.80 2,3	1	$(^3\text{P}_1)5\text{d}$	[2]3/2	132207.76	$(^1\text{D}_2)6\text{p}$	[2]5/2	noise?
4818.02	200	4817.98	-0.04	-1.25	1,2,4	$(^3\text{P}_2)5\text{d}$	[2]3/2	116783.09	$(^3\text{P}_2)6\text{p}$	[1]3/2	3 artifact
4823.35	300	4823.25	-0.10	-0.65	1,2,3	$(^3\text{P}_2)6\text{p}$	[2]3/2	132518.82	$(^3\text{P}_2)7\text{s}$	[2]5/2	4 blend
4844.33	2000	4844.33	0.00	+0.65 +0.60 2,3,4 +0.491N +0.510±0.027ZBD	1	$(^3\text{P}_2)6\text{s}$	[2]5/2	113705.40	$(^3\text{P}_2)6\text{p}$	[3]7/2	
4876.50	500	4876.50	0.00	+0.10 +0.255N	1,2,3	$(^1\text{D}_2)6\text{s}$	[2]5/2	130063.96	$(^1\text{D}_2)6\text{p}$	[3]7/2	4 blend
4883.53	600	4883.53	0.00	-0.25	1,2,3	$(^3\text{P}_0)6\text{s}$	[0]1/2	121628.82	$(^3\text{P}_0)6\text{p}$	[1]3/2	
4884.15	100	4884.09	-0.06	-0.80	1	$(^3\text{P}_0)6\text{p}$	[1]1/2	140883.42	$(^3\text{P}_0)7\text{s}$	[0]1/2	2,3,4 not obs.
4887.30	300	4887.30	0.00	-0.90 -0.80 2,3	1,4	$(^3\text{P}_1)6\text{s}$	[1]3/2	123254.60	$(^3\text{P}_1)6\text{p}$	[2]3/2	
4890.090	300	4890.085	-0.005	-1.20 -1.10 2 -0.754±0.022ZBD	1,3,4	$(^3\text{P}_2)6\text{s}$	[2]5/2	113513.36	$(^3\text{P}_2)6\text{p}$	[3]5/2	
4919.66	200	4919.66	0.00	-0.95 -0.65 -0.85	1,2 3 4	$(^3\text{P}_1)5\text{d}$	[1]1/2	124571.09	$(^3\text{P}_1)6\text{p}$	[1]1/2	
4921.48	800	4921.48	0.00	+0.10 -0.10 4	1,2,3	$(^3\text{P}_1)6\text{s}$	[1]3/2	123112.54	$(^3\text{P}_1)6\text{p}$	[2]5/2	
4971.71	200	4971.68	-0.03	-0.75	1,2,3,4	$(^1\text{D}_2)5\text{d}$	[3]5/2	139193.80	$(^3\text{P}_2)7\text{p}$	[1]3/2	2,3,4 noise
4972.71	400	4972.70	-0.01	-0.55	1,2,3,4	$(^1\text{D}_2)6\text{s}$	[2]5/2	129667.35	$(^1\text{D}_2)6\text{p}$	[1]3/2	
4988.77	300	4988.725	-0.045	-1.00 -0.80 2,3,4	1	$(^3\text{P}_1)5\text{d}$	[1]1/2	124289.45	$(^3\text{P}_1)6\text{p}$	[1]3/2	blend
5044.92	150	5044.92	0.00	-0.80	1,2,4	$(^1\text{D}_2)6\text{s}$	[2]3/2	132741.15	$(^1\text{D}_2)6\text{p}$	[1]1/2	3,4 noise
5080.62	600	5080.51	-0.11	-0.30 -0.05 2	1,3	$(^3\text{P}_2)6\text{p}$	[3]5/2	133189.42	$(^3\text{P}_2)7\text{s}$	[2]3/2	
5122.42	200	5122.31	-0.11	-0.50 -0.30 2,3	1	$(^3\text{P}_2)6\text{p}$	[1]1/2	133189.42	$(^3\text{P}_2)7\text{s}$	[2]3/2	4 noise 3 blend
5188.04	200	5188.08	+0.04	-1.10	1,3	$(^3\text{P}_1)6\text{p}$	[2]5/2	142382.13	$(^3\text{P}_1)7\text{s}$	[1]3/2	2 blend, 4 not obs

**Table B.1.** cont.

$\lambda$ (Lab)	Int.	$\lambda$ (stellar)	$\Delta\lambda$	log gf	$\chi_{low}$ (cm $^{-1}$ )	Term	$\chi_{up}$ (cm $^{-1}$ )	Term	Notes
5260.44	200	5260.42	-0.02	-0.437N -0.25	1,4 2	104250.06 ( $^3P_1$ )5d	[1]1/2	123254.60 ( $^3P_1$ )6p	[2]3/2
		5260.44	0.00	-0.35	3				
5261.95	200	5261.95	0.00	+0.25 +0.150N	1,2,3	112924.84 ( $^1D_2$ )6s	[2]3/2	131923.79 ( $^1D_2$ )6p	[2]3/2
5268.31	50	5268.25	-0.06	-1.00 -0.80 -0.70	1 2 3,4	105313.33 ( $^3P_2$ )5d	[1]3/2	124289.45 ( $^3P_1$ )6p	[1]3/2
5292.22	1000	5292.22	0.00	+0.60 +0.45 +0.351N +0.382±0.013ZBD	1 2,3,4	93068.44 ( $^3P_2$ )6s	[2]5/2	111958.89 ( $^3P_2$ )6p	[2]5/2
5309.27	200	5309.27	0.00	-0.95	1,2,3,4	102799.07 ( $^3P_1$ )6s	[1]3/2	121628.82 ( $^3P_0$ )6p	[1]3/2
5313.87	800	5313.76	-0.11	-0.10 -0.15 -0.05	1 2 3,4	113705.40 ( $^3P_2$ )6p	[3]7/2	132518.82 ( $^3P_2$ )7s	[2]5/2
5339.33	1000	5339.355	+0.025	-0.07 -0.10 -0.15 +0.048±0.019ZBD	1,4 2 3	93068.94 ( $^3P_2$ )6s	[2]5/2	111792.17 ( $^3P_2$ )6p	[2]3/2
5368.07	100	5368.075	+0.005	-1.05	1,2,3	105947.55 ( $^3P_2$ )5d	[0]1/2	124571.09 ( $^3P_1$ )6p	[1]1/2
5372.39	300	5372.405	+0.015	-0.211N -0.10	1,4 2,3	95064.38 ( $^3P_2$ )6s	[2]3/2	113672.89 ( $^3P_2$ )6p	[1]1/2
blend									
5419.15	2000	5419.155	+0.005	+0.42 +0.35 +0.215N +0.256±0.015ZBD	1 2,3,4	95064.38 ( $^3P_2$ )6s	[2]3/2	113512.36 ( $^3P_2$ )6p	[3]5/2
5438.96	400	5438.96	0.00	-0.44 -0.183N	1,2,3,4	102799.07 ( $^3P_1$ )6s	[1]3/2	121179.80 ( $^3P_1$ )6p	[0]1/2
5450.45	100	5450.45	0.00	-1.10 -0.90	1 2,3	105947.55 ( $^3P_2$ )5d	[0]1/2	124289.45 ( $^3P_1$ )6p	[1]3/2
5460.39	300	5460.365	-0.025	-0.85 -0.75 -0.673±0.030ZBD	1 2,3,4	95396.74 ( $^3P_2$ )5d	[2]5/2	113705.40 ( $^3P_2$ )6p	[3]7/2
5472.61	500	5472.60	-0.01	-0.55 -0.449N -0.362±0.030ZBD	1,2,3,4	95437.67 ( $^3P_2$ )5d	[3]7/2	113705.40 ( $^3P_2$ )6p	[3]7/2

**Table B.1.** cont.

$\lambda$ (Lab)	Int.	$\lambda$ (stellar)	$\Delta\lambda$	$\log gf$	$\chi_{low}(\text{cm}^{-1})$	Term	$\chi_{up}(\text{cm}^{-1})$	Term	Notes
5531.07	400	5531.05	-0.02	-0.90 -0.80 -0.616N $-0.632 \pm 0.021 \text{ZBD}$	1 2,3 4	$95437.67$ ( <sup>3</sup> P <sub>2</sub> )5d	[3]7/2 [3]5/2	113512.36 ( <sup>3</sup> P <sub>2</sub> )6p	
5616.67	150	5616.65	-0.02	-0.80 -0.40	1,3,4 2	105313.33 ( <sup>3</sup> P <sub>2</sub> )5d	[1]3/2	123112.54 ( <sup>3</sup> P <sub>1</sub> )6p	[2]5/2 blend
5659.38	150	5659.38	0.00	-0.80 -0.50	1,3 2,4	106906.12 ( <sup>3</sup> P <sub>1</sub> )6s	[1]1/2	124571.09 ( <sup>3</sup> P <sub>1</sub> )6p	[1]1/2 noise ?
5667.56	300	5667.540	-0.02	-0.65 -0.50 -0.45	1 2,3 4	96033.48 ( <sup>3</sup> P <sub>2</sub> )5d	[2]3/2	113672.89 ( <sup>3</sup> P <sub>2</sub> )6p	[1]1/2
5699.61	100	5699.61	0.00	-0.85?	1	111326.96 ( <sup>3</sup> P <sub>1</sub> )5d	[2]3/2	128867.20 ( <sup>1</sup> D <sub>2</sub> )6p	[3]5/2 2,3,4 noise
5719.61	200	5719.587	-0.023	-0.80 $-0.746N$ $-0.687 \pm 0.023 \text{ZBD}$	1,2	96033.48 ( <sup>3</sup> P <sub>2</sub> )5d	[2]3/2	113512.36 ( <sup>3</sup> P <sub>2</sub> )6p	[3]5/2 3,4 blend telluric
5726.91	200	5726.88	-0.03	-0.35 -0.25	1 2,4	114751.08 ( <sup>3</sup> P <sub>2</sub> )5d	[3]5/2	132207.76 ( <sup>1</sup> D <sub>2</sub> )6p	[2]5/2 3 blend telluric
5751.03	200	5750.99	-0.04	-0.35 -0.45	1,2 3,4	106906.12 ( <sup>3</sup> P <sub>1</sub> )6s	[1]1/2	124289.45 ( <sup>3</sup> P <sub>1</sub> )6p	[1]3/2 2 noise
5758.65	100	5758.665	+0.015	-0.35	1,4	112703.64 ( <sup>3</sup> P <sub>1</sub> )5d	[2]5/2	130063.96 ( <sup>1</sup> D <sub>2</sub> )6p	[3]7/2 blend, 2,3 noise
5776.39	100	5776.39	0.00	-0.70	1	105947.55 ( <sup>3</sup> P <sub>2</sub> )5d	[0]1/2	123254.60 ( <sup>3</sup> P <sub>1</sub> )6p	[2]3/2 2,3 not obs, 4 no spectrum
5893.29	150	5893.29	0.00	-0.90	1	112703.64 ( <sup>3</sup> P <sub>1</sub> )5d	[2]5/2	129667.35 ( <sup>1</sup> D <sub>2</sub> )6p	[1]3/2 2 noise, 3 not obs, 4 blend
5905.13	200	5905.115	-0.015	-0.85 -0.65	1 2	104250.06 ( <sup>3</sup> P <sub>1</sub> )5d	[1]1/2	121179.80 ( <sup>3</sup> P <sub>1</sub> )6p	[0]1/2 2,3 blend telluric
5945.53	300	5945.53	0.00	-0.60 -0.80	2,4 3	96958.18 ( <sup>3</sup> P <sub>2</sub> )5d	[1]1/2	113672.89 ( <sup>3</sup> P <sub>2</sub> )6p	[1]1/2 1 blend telluric
5971.13	200	5971.135	+0.005	-0.50	1	112924.84 ( <sup>1</sup> D <sub>2</sub> )6s	[2]3/2	129667.35 ( <sup>1</sup> D <sub>2</sub> )6p	[1]3/2 2,3,4 noise
5976.46	1000	5976.460	+0.00	-0.222N $-0.317 \pm 0.023 \text{ZBD}$	1,2 3,4	95064.38 ( <sup>3</sup> P <sub>2</sub> )6s	[2]3/2	111792.17 ( <sup>3</sup> P <sub>2</sub> )6p	[2]3/2
6036.20	500	6036.170	-0.03	-0.57 -0.45 -0.609N $-0.562 \pm 0.020 \text{ZBD}$	1 2 3,4	95396.74 ( <sup>3</sup> P <sub>2</sub> )5d	[2]5/2	111958.89 ( <sup>3</sup> P <sub>2</sub> )6p	[2]5/2
6051.15	1000	6051.120	-0.03	-0.252N $-0.257 \pm 0.020 \text{ZBD}$	1,2,3 4	95437.67 ( <sup>3</sup> P <sub>2</sub> )5d	[3]7/2	111958.89 ( <sup>3</sup> P <sub>2</sub> )6p	[2]5/2
6097.59	1000	6097.57	-0.02	-0.45 -0.35 -0.30 -0.237N $-0.355 \pm 0.025 \text{ZBD}$	1,3 2 4 3	95396.74 ( <sup>3</sup> P <sub>2</sub> )5d	[2]5/2	111792.17 ( <sup>3</sup> P <sub>2</sub> )6p	[2]3/2
6101.43	200	6101.37	-0.06	-0.70 -0.10	1,4 3	107904.50 ( <sup>3</sup> P <sub>1</sub> )5d	[1]3/2	124289.45 ( <sup>3</sup> P <sub>1</sub> )6p	[1]3/2 2 noise, 4 blend

