


## Exoplanets Direct imaging

Planets and Astrobiology (2018-2019)  
G. Vladilo

1

Black body flux (in units  $10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1}$ ) of some Solar System bodies as “seen” from 10 pc. A putative hot Jupiter is also shown. The planets have two peaks in their spectra. The short-wavelength peak is due to sunlight scattered from the planet using the planet’s albedo. The long-wavelength peak is from the planet thermal emission estimated by a black body of the planets’s effective temperature.

Seager & Deming (2010)



3

## Direct method of exoplanet detection

- **Direct imaging**
  - The image of the planet is searched for in the field of the star
- **Observational challenges**
  - Planet-star luminosity contrast
  - Planet-star angular separation
- **Both challenges set strong constraints on the observational techniques**
  - The luminosity contrast can be quantified with simulations of stellar and planet spectra

2

## Direct imaging: observational challenges

- **Luminosity contrast**

$$L_p/L_*$$

- Optical spectral band  
reflected stellar radiation  
varies with the orbital phase  
contrast  $\sim 10^9$ - $10^{10}$

- Infrared spectral band ( $\sim 10 \mu\text{m}$ )  
intrinsic planetary emission  
contrast  $\sim 10^6$ - $10^7$

Optical band

$$L_p \sim L_* \left( \frac{R_p}{a} \right)^2 \Phi(t)$$

$$\Phi(t) = 1 - \sin i \sin \left( \frac{2\pi t}{P} \right)$$


Infrared band

$$L_p \sim L_* (M_p/M_*)$$

4

### Luminosity contrast versus wavelength


Models for giant planets orbiting a solar-type star, cloud effects included



5

### Luminosity contrast versus wavelength


Models for giant planets orbiting a solar-type star, cloud effects included



7

### Luminosity contrast versus wavelength

Models for giant planets orbiting a solar-type star, cloud effects included



6

### Direct imaging: observational challenges

- Planet-star angular separation

Angular separations can be estimated as a function of stellar distance,  $l$ , and orbital semimajor axis of the planet,  $a$

$$\vartheta = \arctan \frac{a}{l}$$

Typical values are lower than 1 arcsec

e.g., the Earth-Sun separation as seen from 20 pc is 50 mas

The luminosity contrast makes hard to attain the theoretical diffraction limit  $\delta\vartheta \approx \lambda/D$  (radius of the Airy disk)

$D$ : telescope diameter


$\lambda$ : wavelength of the observations

8

## Angular separation and luminosity contrast

The maximum luminosity contrast increases with angular separation

Dashed line: example of maximum luminosity contrast that can be attained for a given instrumental setup



9

## Direct imaging of exoplanets

- **Observable quantities**

- Orbital parameters

Given the stellar distance  $l$ , one can determine the orbital semimajor axis  $a$  from the angular separation  $\vartheta$

Given  $a$ , the orbital period  $P$  is estimated with the third Kepler's law

- Effective temperature

The effective temperature can be determined from the energy distribution of the planet spectrum

The spectral distribution can be estimated by comparing photometric measurements taken in different spectral bands

- Emission spectrum of the planet

If the source is sufficiently bright, a spectrum can be taken

The spectrum can be used to study the (atmospheric) chemical composition of the planet

11

## Direct imaging of exoplanets

- **Observational bias**

- Due to the observational challenges specific of the direct imaging method, the following types exoplanets are preferentially found

- (1) around nearby stars (small  $l$ )

- (2) with wide orbits (large  $a$ )

- (3) with high intrinsic emission (e.g. young and massive)

- (4) with high planet/star mass ratio  $M_p/M_*$

10

## Direct imaging of exoplanets

- **Model-dependent planetary parameters**

- Planet mass and size

Mass and size can be estimated combining the measurement of planet emissivity with a model of planet evolution

The error on the masses estimated in this way can be quite large and is also related to the uncertain age of the system

- **Advantages of the direct imaging method**

- Allows us to study planets distant from the star

- Not affected by the temporal baseline bias (a single observation yields  $a$ , from which one can determine  $P$ )

- Not affected by variability of the central star

- Best way to obtain *direct* informations on the properties of exoplanets

12

**Direct imaging of exoplanets:**  
Techniques to deal with the luminosity contrast

• **Coronagraphic techniques**

- High contrast can be achieved by rejecting the stellar light from the area of interest in the focal plane
- The technique, employing some form of mask in the telescope focal plane, is referred to as *coronagraphy*

Coronagraphy was originally developed to study solar corona, a tenuous structure of highly ionized gas that surrounds the Sun, which is hard to observe due to the extremely high luminosity contrast with the solar disk

Ideally, a coronagraph coupled to an adaptive optics system would perform as if placed above the atmosphere

**Direct imaging of exoplanets:**  
Techniques to increase the angular resolution

Ground-based telescopes do not attain the theoretical diffraction limit because the resolution is limited by the turbulence in the atmosphere


The atmosphere has density enhancements (turbulent cells) of size  $d_0 \sim 0.1\text{m}$  that are carried across the telescope line of sight by high-altitude winds  
 Segments of the wavefront comparable to the size of the turbulent cells will be nearly planar; they are called isophase patches  
 Each isophase patch uses  $\sim 0.1\text{m}$  of the telescope aperture and its image is the Airy disk of a  $0.1\text{m}$  telescope, i.e.  $\lambda d_0 \sim 1''$   
 The summation of the images from all the isophase patches across the line of sight will yield an image with size  $\sim 1''$ , much larger than the theoretical angular resolution  $\delta\theta \sim \lambda/D$

**Direct imaging of exoplanets:**  
Techniques to deal with the luminosity contrast

• **Developments and limits of coronagraphic techniques**

- Alternative concepts for high-rejection coronagraphs have been stimulated by the interest in exoplanet imaging
- A variety of designs theoretically able to achieve  $10^{10}$  contrast within  $5 \lambda/D$  are being developed

Throughput at the  $10^{10}$  contrast level




**Techniques to deal with atmospheric turbulence:**  
Adaptive optics

Consists in the real time control of the optics of a telescope to counteract atmospheric turbulence  
 Adaptive optic systems must be very fast since atmospheric turbulence varies with a time scale of order  $\sim 1$  millisecond  
 Adaptive optics requires an ancillary system that senses the instantaneous shape of the atmospheric wavefront  
 Wavefront sensing is performed using a bright reference star and/or the atmospheric reflection of a laser beam which acts as an artificial star  
 The information on the wavefront is sent to hardware and analysis software which detects deviations from a planar wave with proper centroid location  
 Corrections are then applied to deformable mirrors located along the optical path; the modified beam is then sensed and the process continues in this feedback-loop mode

## Sketch of a complete adaptive optics system

The loop must work on time scales  $\leq 1$  ms

The movable flat mirror moves the image centroid to keep it in a single position in the focal plane; this reduces the excursions required on the actuators of the deformable mirror



17

## • Speckle imaging techniques


- In the *image stacking*, the short exposure images are lined up by the brightest speckle and averaged together to give a single image
- In the *Lucky Imaging* approach, only the best few short exposures are selected
- *Speckle interferometry* makes use of Fourier analysis to obtain the high-resolution structure of the object from the speckle patterns
- Practical procedure
  - Record many frames rapidly
  - Take the power spectrum of each frame and average
  - Divide the power spectrum of the target by the power spectrum of a point source
  - Fit a model (e.g. planet around a central star) to the true Fourier transform of the object

19

## Techniques to deal with atmospheric turbulence: Speckle interferometry

It works by obtaining images of the object sufficiently rapidly (e.g.,  $\sim 1$ -10 ms) to freeze the blurring that arises from atmospheric scintillation

The resulting image of a point source then consists of a large number of small dots or speckles, each of which is a diffraction limited image with effective value of  $d$  up to the actual diameter of the telescope




Example: speckle images of Vega first obtained in the '70s

18

## Direct imaging of exoplanets Examples

- Fomalhaut b
  - discovered in 2008 from a re-analysis of previous HST data
  - $a = 119$  AU;  $e \sim 0.11$ ;  $M \sim 3 M_J$ ;  $P \sim 870$  yr
  - Fomalhaut:  $d = 8$  pc
  - Kalas et al. (2008)



20

## Direct imaging of exoplanets

### Examples

AB Pic b

$$M = 13 M_J$$

close to the brown dwarf boundary

$$a > 80 \text{ AU}$$


AB Pic, K2 V

$$d = 47 \text{ pc}$$

Discovered using near-infrared Lyot coronagraphic observations

NACO at ESO-VLT

Chauvin et al. (2005)



21

## Multiple planetary systems detected with direct imaging

Example:

HR 8799 b, c, d


$$M = 7, 10, 10 M_J$$

$$a = 68, 38, 24 \text{ AU}$$

$$d = 39 \text{ pc}$$

Coronagraphic and speckle imaging techniques in the infrared

(Marois et al. 2008)



## Direct imaging of exoplanets

### Examples

GJ 504 b

$$M = 4 M_J$$

$$a = 44 \text{ AU}$$


Sun-like star GJ 504

$$d = 17.6 \text{ pc}$$

Lowest temperature (510 K) exoplanet observed with direct imaging

Adaptive optics, occulting mask, near infrared

Kuzuhara et al. (2013)




22

## Direct imaging: summary of results

About a hundred planets has been detected with the direct imaging method. These planets are quite massive and distant from the central star.

Mass – Period Distribution



24


## Direct imaging of exoplanets Future perspectives

- Occulters
- Infrared space interferometers
- Nulling interferometry
- Space projects (suspended)
  - ESA Darwin
  - NASA Terrestrial Planet Finder (TPF)
- Ground based projects
  - examples:
    - SPHERE at ESO VLT (active)
    - EPICS at ESO E-ELT (future development)



25

- Nulling interferometry
  - Introduces destructive interference between the pupils of two telescopes for an on-axis star
  - Identical path lengths through the two beams leads to an interference maximum for an on-axis source
  - Introducing a phase difference of  $\pi$  rad in one of the paths suppresses the central maximum
  - By varying the baseline  $D$ , a range of constructive interference angles can be examined for the presence of an off-axis source



26