

Climate and habitability

Planets and Astrobiology (2018-2019)
G. Vladilo

1

Complexity of the climate system: the problem of the time scales

Time scales of different components of the climate system

<i>Atmosphere</i>		
Overall response time to heating		months
Typical spin-down time of wind if nothing is forcing it		days
Frontal system lifetime (1000s of km)		days
Convective cloud lifetime (100 m to km horizontal; up to 10 km vertical)		hours
Time scale for typical upper-level wind (20 m s^{-1}) to cross continent (a few 1000 km)		days
<i>Ocean</i>		
Response time of upper ocean (above thermocline) to heating		months to years
Response time of deep ocean to atmospheric changes		decades to millennia
Ocean eddy lifetime (10s to 100 km)		months
Ocean mixing in the surface layer		hours to days
Time for typical ocean current (cm s^{-1}) to cross ocean (1000s of km)		decades
<i>Cryosphere</i>		
Snow cover		months
Sea ice (extent and thickness variations)		months to years
Glaciers		decades to centuries
Ice caps		centuries to millennia
<i>Land surface</i>		
Response time to heating		hours
Response time of vegetation to oppose excess evaporation		hours
Soil moisture response time		days to months
<i>Biosphere</i>		
Ocean plankton response to nutrient changes		weeks
Recovery time from deforestation		years to decades
<i>Lithosphere</i>		
Isostatic rebound of continents (after being depressed by weight of glacier)		10 000s of years
Weathering, mountain building		1 000 000s of years

3

Climate models and planetary habitability


- The calculation of the physical conditions at the planetary surface requires the use of climate models
 - With climate models we can take into account the greenhouse effect and a variety of other processes that affect the surface planetary conditions
- Climate models, originally developed for Earth studies, are becoming a key tool for modeling planetary habitability
 - The state-of-the-art models, called “Global Circulation Models” (GCM), are extremely time consuming
 - Simplified climate models are often used for studies of planetary habitability

Bibliographic material:

Pierrehumbert (2010) Principles of Planetary Climate
Neelin (2011) Climate Change and Climate Modeling

2


Reductionistic approaches to tackle the complexity of the climate system



Provenzale (2013)

4

Reductionistic approaches to tackle the complexity of the climate system




Provenzale (2013)

5

Climate instabilities


can drive the planet temperature out of the range of habitability

- **The ice-albedo feedback**
 - If water and ice are present on the planet, a decrease of temperature will increase the surface covered by ice
 - Ice has a very high albedo and an increase of the ice extension will cool the planet even more
 - In extreme conditions, this feedback may lead to a “snowball planet”, i.e. a planet fully covered by ice



7

Reductionistic approaches to tackle the complexity of the climate system




Provenzale (2013)

6

Climate instabilities

can drive the planet temperature out of the range of habitability

- **The temperature-water vapour feedback**
 - If water is present on the planet, the water vapour pressure rises with temperature
 - Water vapour is a strong greenhouse gas, and a rise of water vapour will rise the temperature even more
 - In normal conditions, this feedback is not catastrophic because the cooling rate scales as σT^4



8

The classic habitable zone

- Early calculations of planetary habitability were performed before exoplanets were discovered
 - J. Kasting and collaborators (Penn State University)
- Simplified climate models
 - Radiative-convective transport in a single atmospheric column
- Calculated for stars of different spectral types
 - The energy distribution of the stellar spectrum affects the albedo
- Definition of the circumstellar habitable zone
 - Interval of distances from the central star where a habitable planet can be found
 - Different types of criteria are adopted to define the inner and outer edge of the habitable zone
 - Climate instabilities play an important role in the definition of the habitable zone


9

The inner edge of the habitable zone

- The runaway greenhouse mechanism
 - If the temperature-water vapour feedback is extreme, the vapour may reach the outer layers of the atmosphere
 - In the outer layers the water molecules can be dissociated by high energy stellar photons
 - The hydrogen produced by photodissociation can be lost to space
 - This chain of events is called the runaway greenhouse mechanism
 - In the long term, this mechanism may lead to the disappearance of liquid water on the planet
 - The “runaway greenhouse” mechanism is used to define the inner edge of the habitable zone

11

The classic habitable zone



10

The outer edge of the habitable zone

- An increase of greenhouse gases in the planetary atmosphere makes the planet habitable at lower levels of stellar flux, i.e. at larger distances from the central star
- The outer edge of the habitable zone is defined assuming that the planetary atmosphere is dominated by CO_2
 - as in the case of Mars
- The amount of CO_2 that is able to warm the planet at low levels of insolation is limited by the onset of CO_2 clouds with high albedo, which would counteract the heating due to greenhouse effect

12

Mechanisms of climate stabilization

In the definition of the classic habitable zone it is assumed that the planet has the capability of adjusting the level of CO_2 through a mechanism of climate stabilization

The fact that Earth's climate has been relatively stable in the course of geological time scales suggests the existence of a mechanism of climate stabilization

This mechanism must have been able to stabilize the Earth's climate despite changes that have occurred in terms of solar radiation, atmospheric composition and other factors

The mechanism invoked for the Earth is based on a CO_2 inorganic cycle

13

The CO_2 cycle of climate stabilization

The weathering efficiency increases with atmospheric temperature


The rate of CO_2 emission is independent of the atmospheric temperature

As a result, there is a negative feedback temperature- CO_2 that stabilizes the climate

- The time scale of the cycle is estimated to be $\sim 5 \times 10^5$ years


15

The CO_2 cycle of climate stabilization



14

- The existence of tectonics and volcanism is necessary for the existence of the CO_2 stabilization cycle
 - In the present-day Solar System, only the Earth has these types of geophysical activities
 - The CO_2 cycle of climate stabilization is invoked in the definition of the classic habitable zone
 - How common this mechanism can be in terrestrial-type exoplanets is currently the subject of studies of planetary interiors




16

Habitability under the planet surface

- The definition of habitable zone relies on the concept of surface habitability
 - Habitability under the planet surface could be present in planetary bodies outside the circumstellar habitable zone, in particular beyond the outer edge
- Temperature and pressure gradients may yield conditions of habitability in the interior of planets or satellites
 - Internal sources of heat yield a temperature gradient in the planet interior
 - The pressure gradient towards the planetary interior may improve the conditions of habitability

17

Energy balance models (EBM) of planetary climate




$$I_i + C_i \frac{\partial T}{\partial t} - \frac{\partial}{\partial x} \left[D_i (1 - x^2) \frac{\partial T}{\partial x} \right] = S_i (1 - A_i)$$

$$x = \sin \phi \quad (\phi \text{ is the latitude})$$

19

Surface habitability with Energy Balance Models (EBMs)

Simplified climate models aimed at predicting the seasonal and latitudinal distribution of the surface temperature



18

Brief summary of recent studies of planetary habitability carried out at INAF-OATs

THE ASTROPHYSICAL JOURNAL, 767:65 (23pp), 2013 April 10
 © 2013. The American Astronomical Society. All rights reserved. Printed in the U.S.A. doi:10.1088/0004-637X/767/1/65

THE HABITABLE ZONE OF EARTH-LIKE PLANETS WITH DIFFERENT LEVELS OF ATMOSPHERIC PRESSURE

GIOVANNI VLADILLO^{1,2}, GIUSEPPE MURANTE¹, LAURA SILVA¹, ANTONELLO PROVENZALE³, GAIA FERRI², AND GREGORIO RAGAZZINI²

THE ASTROPHYSICAL JOURNAL, 804:50 (20pp), 2015 May 1
 The American Astronomical Society. All rights reserved. doi:10.1088/0004-637X/804/1/50

MODELING THE SURFACE TEMPERATURE OF EARTH-LIKE PLANETS

GIOVANNI VLADILLO^{1,2}, LAURA SILVA¹, GIUSEPPE MURANTE^{1,3}, LUCA FILIPPI^{3,4}, AND ANTONELLO PROVENZALE^{3,5}

¹INAF—Trieste Astronomical Observatory, Trieste, Italy; vladilo@oats.inaf.it
²Department of Physics, University of Trieste, Trieste, Italy
³Institute of Atmospheric Sciences and Climate—CNR, Torino, Italy
⁴DIMEAS, Politecnico di Torino, Torino, Italy
⁵Institute of Cosmosciences and Earth Resources—CNR, Pisa, Italy
 Received 2015 January 29; accepted 2015 April 3; published 2015 April 29


From climate models to planetary habitability: temperature constraints for complex life

Laura Silva¹, Giovanni Vladilo¹, Patricia M. Schulte², Giuseppe Murante¹, Antonello Provenzale³

20

Examples of application of the ESTM
Seasonal and latitudinal surface temperature of the Earth


Variation of stellar insolation



21

Examples of application of the ESTM
Seasonal and latitudinal surface temperature of an Earth-like planet


Variation of rotation period



23

Examples of application of the ESTM
Seasonal and latitudinal surface temperature of the Earth

Variation of surface pressure



22

Quantifying the habitability with
the liquid water criterion

The climate simulation yields
the surface temperature
as a function of latitude and time

$$T(\phi, t)$$

Habitability function $H(\phi, t) = \begin{cases} 1 & \text{if } T_{\text{melt}}(p) \leq T(\phi, t) \leq T_{\text{boil}}(p) \\ 0 & \text{otherwise} \end{cases}$


Liquid water
criterion

Mean global
annual
habitability

$$h = \frac{\int_{-\pi/2}^{+\pi/2} d\phi \int_0^P dt [H(\phi, t) \cos \phi]}{2P}$$


24

At low pressure,
the mean planet temperature is not a good diagnostic of habitability




Evaporating planets inside the "habitable zone"

Effects of eccentricity



Pressure dependence of the habitable zone
Obtained running a large number of climate simulations
Vladilo et al. (2013)



Self-consistent treatment of surface temperature and pressure in the definition of habitability by means of the liquid water criterion

The atmospheric mass habitable zone
for complex life and atmospheric biomarkers
Obtained using the temperature limits $0^{\circ}\text{C} \leq T \leq 50^{\circ}\text{C}$
Silva et al. (2017)

