

# Astrobiology

## Lecture 3


### Water & carbon in terrestrial life

Trieste University, Academic Year 2021-2022  
Giovanni Vladilo (INAF-OATs)


# The water molecule

Builds up the molecular medium of terrestrial life

The water molecule is polar



(a) Electron shells in a water molecule



(b) Distribution of partial charges in a water molecule

## Polar and non-polar molecules


- The polar character depends on the geometrical distribution of electric charges of the molecule

Water is polar because of the asymmetric distribution of charges



Methane is non polar (no electric dipole)

Methane:  
a non-polar molecule

- Polar molecules
  - can be solved in water
  - are hydrophilic
- Non-polar molecules
  - cannot be solved in water
  - are hydrophobic



Water molecules are connected by a network of hydrogen bonds




Hydrogen bonds are responsible for most of the properties of water that are relevant for life

# Properties of water relevant to life (1)

- The water molecule has a high electric dipole
  - Water is a good solvent

Thanks to this property, life molecular constituents can be dissolved and have the mobility required for metabolic processes to take place
  - Thanks to the polarity, amphiphilic molecules in water can spontaneously form structures of biological interest (bilayers, micelles)

Polar molecules allow spontaneous formation  
of molecular structures of biological interest



## Properties of water relevant to life (2)


- Water spontaneously form ions
  - Spontaneous breaking of covalent bonds in a small fraction of water molecules yields  $\text{H}^+$  and  $\text{OH}^-$  ions

Note: the concentration of  $\text{H}^+$  ions in water is used to define the pH scale

The free ions, and in particular  $\text{H}^+$ ,  
can be used to transport electric charges

$\text{H}^+$  and  $\text{OH}^-$  take part in metabolic reactions

- Water takes part of fundamental metabolic processes, both as a reactant and as a product of reaction
- Water formation and dissociation plays a direct role in metabolic processes





# Carbon in terrestrial life


Electronic orbitals

Carbon-based biological macro-molecules

# Carbon

In terrestrial life carbon is the building block of biological molecules

- With respect to other cosmically abundant atoms, carbon offers several advantages in terms of structural and metabolic properties
- Electronic configuration
  - Carbon's ground state configuration is  $1s^2 2s^2 2p^2$




- The excitation of one electron of the 2s orbital easily provides a configuration with 4 orbitals with a single electron
- The 4 oriented covalent bonds allow the formation of a great variety of 3D molecular structures:

linear, planar, tetrahedral


# Hybridization of carbon valence orbitals

**sp orbitals:** two atomic orbitals are mixed to form two hybrid orbitals





The two sp hybrid orbitals arrange themselves in three dimensional space to get as far apart as possible with a bond angle of  $180^\circ$ .

The geometry which achieves is linear.





## Hybridization of carbon valence orbitals

**$sp^2$  orbitals:** three atomic orbitals are mixed to form three hybrid orbitals



**$sp^3$  orbitals:** four atomic orbitals are mixed to form four hybrid orbitals




# Carbon

- **Structural properties**
  - In summary, carbon has 4 oriented covalent bonds that allow the formation of a great variety of 3D molecular structures
  - The valence orbitals 2s and 2p can hybridize forming:
    - two sp hybrid orbitals → linear structures
    - three sp<sup>2</sup> hybrid orbitals → planar structures
    - four sp<sup>3</sup> hybrid orbitals → tetrahedral structures
  - The same flexibility of forming geometrical structures is not found in other atoms


## Carbon versus nitrogen and oxygen

The flexibility of carbon to form 3D structures is not found in other atoms

Nitrogen has 3 covalent bonds which tend to form planar structures




Oxygen has 2 covalent bonds which tend to form linear structures




## Organic ring structures

Carbon can form a variety of ring structures

Carbon is the only atom with the capability of forming aromatic rings



Cyclohexane  
(saturated molecule)




Aromatic ring of Benzene  
(unsaturated molecule)

# Hetero-organic molecules

- Carbon is capable of forming complex molecules not only with itself, but also with H, O and N

This is because the bonds C-C, C-H, C-O, and C-N have similar energies

For instance, N can replace C in ring structures



The large flexibility of carbon in terms of geometrical structure, coupled with the possibility of substitutions of other abundant elements, leads to a infinite number of possible molecular structures potentially suitable for different biological functions



# Advantages of carbon

## Metabolic properties

- Carbon can easily be transformed from the completely oxidized form,  $\text{CO}_2$ , to the completely reduced form,  $\text{CH}_4$

This is an advantage for the capability of activating metabolic processes, which are largely based on redox reactions

This provides the possibility of cycling carbon between its “inorganic form” and its “organic form”

$\text{CO}_2$  : inorganic carbon

$\text{CH}_4$  : organic carbon

The great capability of carbon to form complex structures is supported by astronomical observations of interstellar molecules

All interstellar molecules with at least 6 atoms are organic

**Table 5.4** Some complex carbon compounds detected in the interstellar medium and meteorites.

Number of Atoms				
6	7	8	9	> 9
C <sub>5</sub> H, HCH <sub>2</sub> OH	CH <sub>3</sub> C <sub>2</sub> H	CH <sub>3</sub> OCHO	(CH <sub>3</sub> ) <sub>2</sub> O	(CH <sub>3</sub> ) <sub>2</sub> CO
NH <sub>2</sub> CHO,	CH <sub>3</sub> CHO	CH <sub>3</sub> C <sub>3</sub> N	CH <sub>3</sub> CH <sub>2</sub> OH	HC <sub>9</sub> N
CH <sub>3</sub> CN	HC <sub>5</sub> N, C <sub>6</sub> H	C <sub>7</sub> H, H <sub>2</sub> C <sub>6</sub>	CH <sub>3</sub> CH <sub>2</sub> CN	HC <sub>11</sub> N
CH <sub>3</sub> NC,	CH <sub>3</sub> NH <sub>2</sub>		HC <sub>7</sub> N	C <sub>6</sub> H <sub>6</sub> , C <sub>60</sub> <sup>+</sup>
CH <sub>3</sub> SH	CH <sub>2</sub> CHCN		CH <sub>3</sub> C <sub>4</sub> H, C <sub>8</sub> H	PAHs,
H <sub>2</sub> C <sub>4</sub> ,	C <sub>2</sub> H <sub>4</sub> O		CH <sub>3</sub> C <sub>4</sub> N	glycine?
HCC <sub>2</sub> HO,				
C <sub>5</sub> H, C <sub>5</sub> N,				
C <sub>5</sub> O				

# From carbon atoms to biological macromolecules


Carbon is the most abundant constituent  
of the macromolecules of terrestrial life

# Biological macromolecules

The most important biomolecules of terrestrial life are macromolecules with a large number of atomic units (e.g.,  $10^6 - 10^8$  amu)



Macromolecules result from the polymerization of a large number of subunits (monomers)

Terrestrial life features 4 types of macromolecules:  
carbohydrates, lipids, proteins and nucleic acids



# Biological macromolecules

## Carbohydrates (saccharides)



The most abundant molecules in the biological world

Primary source of chemical energy for most organisms

General formula:  $C_x(H_2O)_y$

Monosaccharides (simple sugars)

Oligosaccharides

From 2 to 10 units of monosaccharides

Polysaccharides

More than 10 monosaccharides

# Biological macromolecules


## Carbohydrates

### Ribose

Simple sugar that is found in RNA

In biology only appears in the form D-ribose


Example of homochirality of biological molecules




# Biological macromolecules

## Lipids


(a) Palmitic acid



(b) Stearic acid



(c) Oleic acid



Heterogeneous class of organic molecules with common solubility properties

Insoluble in water

Soluble in certain types of non-polar solvents

Larger number of C–H bonds with respect to carbohydrates



Used for long-term storage of energy



# Phospholipids

Examples of amphiphilic molecules with a hydrophilic end and a hydrophobic end

In liquid water phospholipids spontaneously form a double layer of molecules (bilayer), with the hydrophobic ends facing each other in the inner part, and the hydrophilic ends facing the water





## Phospholipids and cell membranes

Bilayers of phospholipids are the main structural components of cell membranes

Specialized proteins embedded in the bilayer provide the possibility to exchange molecular constituents in and out of the cell

