

Exoplanets


Direct imaging

Planets and Astrobiology (2022-2023)
Giovanni Vladilo & Paolo Simonetti

Direct method of exoplanet detection

- **Direct imaging**
 - The image of the planet is searched for in the field of the star
- **Observational challenges**
 - Planet-star luminosity contrast
 - Planet-star angular separation
- **Both challenges set strong constraints on the observational techniques**
 - The luminosity contrast can be quantified with simulations of stellar and planet spectra

Black body flux (in units $10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1}$) of some Solar System bodies as “seen” from 10 pc. A putative hot Jupiter is also shown. The planets have two peaks in their spectra. The short-wavelength peak is due to sunlight scattered from the planet using the planet’s albedo. The long-wavelength peak is from the planet thermal emission estimated by a black body of the planets’ effective temperature.



Seager & Deming (2010)

Direct imaging: observational challenges

- **Luminosity contrast**

$$L_p/L_*$$

- Optical spectral band
reflected stellar radiation
varies with the orbital phase
contrast $\sim 10^9$ - 10^{10}

- Infrared spectral band ($\sim 10 \mu\text{m}$)
intrinsic planetary emission
contrast $\sim 10^6$ - 10^7

Optical band

$$L_p \sim L_* \left(\frac{R_p}{a} \right)^2 \Phi(t)$$

$$\Phi(t) = 1 - \sin i \sin \left(\frac{2\pi t}{P} \right)$$


Infrared band

$$L_p \sim L_* (M_p/M_*)$$

Luminosity contrast versus wavelength

Models for giant planets orbiting a solar-type star, cloud effects included


Dependence on orbital distance



Luminosity contrast versus wavelength

Models for giant planets orbiting a solar-type star, cloud effects included

Dependence on age




Planet with 1 M_J at 4 AU

Luminosity contrast versus wavelength

Models for giant planets orbiting a solar-type star, cloud effects included

Dependence on planet mass



Direct imaging: observational challenges

- Planet-star angular separation

Angular separations can be estimated as a function of stellar distance, l , and orbital semimajor axis of the planet, a

$$\vartheta = \arctan \frac{a}{l}$$

Typical values are lower than 1 arcsec

e.g., the Earth-Sun separation as seen from 20 pc is 50 mas

The luminosity contrast makes hard to attain the theoretical diffraction limit $\delta\vartheta \cong \lambda/D$ (radius of the Airy disk)

D : telescope diameter


λ : wavelength of the observations

Also the atmosphere disturbs the observation: best “seeing disk” from the surface is ~ 300 - 400 mas large

Angular separation and luminosity contrast

The maximum luminosity contrast increases with angular separation

Dashed line: example of maximum luminosity contrast that can be attained for a given instrumental setup



Direct imaging of exoplanets

- **Observational bias**
 - Due to the observational challenges specific of the direct imaging method, the following types exoplanets are preferentially found
 - (1) around nearby stars (small l)
 - (2) with wide orbits (large a)
 - (3) with high intrinsic emission (e.g. young and massive) with respect to the central star

Direct imaging of exoplanets

- **Observable quantities**

- Orbital parameters

- Given the stellar distance l , one can determine the orbital semimajor axis a from the angular separation ϑ

- Given a , the orbital period P is estimated with the third Kepler's law

- Effective temperature

- The effective temperature can be determined from the energy distribution of the planet spectrum

- The spectral distribution can be estimated by comparing photometric measurements taken in different spectral bands

- Emission spectrum of the planet

- If the source is sufficiently bright, a spectrum can be taken

- The spectrum can be used to study the (atmospheric) chemical composition of the planet

Direct imaging of exoplanets

- **Model-dependent planetary parameters**

- Planet mass and size

Mass and size can be estimated combining the measurement of planet emissivity with a model of planet evolution

The error on the masses estimated in this way can be quite large and is also related to the uncertain age of the system

- **Advantages of the direct imaging method**

- Allows us to study planets distant from the star
- Not affected by the temporal baseline bias (a single observation yields a , from which one can determine P)
- Not affected by variability of the central star
- Best way to obtain *direct* informations on the properties of exoplanets

Direct imaging of exoplanets:

Techniques to deal with the luminosity contrast

- Coronagraphic techniques

- High contrast can be achieved by rejecting the stellar light from the area of interest in the focal plane
- The technique, employing some form of mask in the telescope focal plane, is referred to as *coronagraphy*


Coronagraphy was originally developed to study solar corona, a tenuous structure of highly ionized gas that surrounds the Sun, which is hard to observe due to the extremely high luminosity contrast with the solar disk

Ideally, a coronagraph coupled to an adaptive optics system would perform as if placed above the atmosphere

Direct imaging of exoplanets: Techniques to deal with the luminosity contrast

- **Developments and limits of coronagraphic techniques**
 - Alternative concepts for high-rejection coronagraphs have been stimulated by the interest in exoplanet imaging
 - A variety of designs theoretically able to achieve 10^{10} contrast within $5 \lambda/D$ are being developed

Throughput
at the 10^{10}
contrast level



Direct imaging of exoplanets: Techniques to increase the angular resolution

Ground-based telescopes do not attain the theoretical diffraction limit because the resolution is limited by the turbulence in the atmosphere

The atmosphere has density enhancements (turbulent cells) of size $d_o \sim 0.1\text{m}$ that are carried across the telescope line of sight by high-altitude winds

Segments of the wavefront comparable to the size of the turbulent cells will be nearly planar; they are called isophase patches

Each isophase patch uses $\sim 0.1\text{m}$ of the telescope aperture and its image is the Airy disk of a 0.1m telescope, i.e. $\lambda/d_o \sim 1''$

The summation of the images from all the isophase patches across the line of sight will yield an image with size $\sim 1''$, much larger than the theoretical angular resolution $\delta\vartheta \sim \lambda/D$

Techniques to deal with atmospheric turbulence: Adaptive optics

Consists in the real time control of the optics of a telescope to counteract atmospheric turbulence

Adaptive optic systems must be very fast since atmospheric turbulence varies with a time scale of order ~ 1 millisecc

Adaptive optics requires an ancillary system that senses the instantaneous shape of the atmospheric wavefront

Wavefront sensing is performed using a bright reference star and/or the atmospheric reflection of a laser beam which acts as an artificial star


The information on the wavefront is sent to hardware and analysis software which detects deviations from a planar wave with proper centroid location

Corrections are then applied to deformable mirrors located along the optical path; the modified beam is then sensed and the process continues in this feedback-loop mode

Techniques to deal with atmospheric turbulence: Speckle interferometry

It works by obtaining images of the object sufficiently rapidly (e.g., $\sim 1-10$ ms) to freeze the blurring that arises from atmospheric scintillation

The resulting image of a point source then consists of a large number of small dots or speckles, each of which is a diffraction limited image with effective value of d up to the actual diameter of the telescope



Example: speckle images of Vega first obtained in the '70s

- Speckle imaging techniques

- In the *image stacking*, the short exposure images are lined up by the brightest speckle and averaged together to give a single image
- In the *Lucky Imaging* approach, only the best few short exposures are selected
- *Speckle interferometry* makes use of Fourier analysis to obtain the high-resolution structure of the object from the speckle patterns
- Practical procedure
 - Record many frames rapidly
 - Take the power spectrum of each frame and average
 - Divide the power spectrum of the target by the power spectrum of a point source
 - Fit a model (e.g. planet around a central star) to the true Fourier transform of the object

Direct imaging of exoplanets

Examples


– Fomalhaut b

discovered in 2008 from a re-analysis of previous HST data

$a = 119 \text{ AU}$; $e \sim 0.11$; $M \sim 3 M_J$; $P \sim 870 \text{ yr}$

Fomalhaut: $d = 8 \text{ pc}$

Kalas et al. (2008)



Direct imaging of exoplanets

Examples

AB Pic b

$$M = 13 M_J$$

close to the brown dwarf boundary

$$a > 80 \text{ AU}$$


AB Pic, K2 V

$$d = 47 \text{ pc}$$

Discovered using near-infrared Lyot coronagraphic observations

NACO at ESO-VLT

Chauvin et al. (2005)



Direct imaging of exoplanets

Examples

GJ 504 b

$$M = 4 M_J$$

$$a = 44 \text{ AU}$$

Sun-like star GJ 504


$$d = 17.6 \text{ pc}$$

Lowest temperature (510 K)

exoplanet observed with direct imaging

Adaptive optics, occulting mask, near infrared

Kuzuhara et al. (2013)



Multiple planetary systems detected with direct imaging

Example:

HR 8799 b, c, d


$M = 7, 10, 10 M_J$

$a = 68, 38, 24 \text{ AU}$

$d = 39 \text{ pc}$

Coronagraphic and speckle imaging techniques in the infrared


(Marois et al. 2008)



Direct imaging: summary of results

About a hundred planets has been detected with the direct imaging method. These planets are quite massive and distant from the central star.

Mass – Period Distribution



Direct imaging: future developments

- **Nulling interferometry**

- Introduces destructive interference between the pupils of two telescopes for an on-axis star
- Identical path lengths through the two beams leads to an interference maximum for an on-axis source
- Introducing a phase difference of π rad in one of the paths suppresses the central maximum
- By varying the baseline D , a range of constructive interference angles can be examined for the presence of an off-axis source

