

Astrobiology

Lecture 10

From molecular replicators to protocells

University of Trieste, 2023
Giovanni Vladilo (INAF-OATs)

Origin of molecular replication and metabolism


- Conceptual “chicken-egg” problem
 - In present-day cells, nucleic acids and proteins are responsible for genetic and metabolic functions, respectively
 - The assemblage of proteins requires the instructions stored in nucleic acids
 - The assemblage of nucleic acids requires the catalytic properties of proteins
 - Each one of these two types of macromolecules requires the previous existence of the other one
- Who came first?
 - Proteins or nucleic acids ?
 - Replication/genetic or metabolic functions ?

Origin of molecular replication and metabolism


- In the past, two opposite schools
 - “Metabolism first”
Emphasizing the initial role of metabolic reactions and the catalytic properties of proteins
 - “Genes first”
Emphasizing the initial role of the genetic system and the capability of nucleic acids to store and transmit information
- Present-day approach
 - Appearance of macromolecules that show both genetic and catalytic properties

The “RNA world”

- Present-day, main stream theory in studies on life’s origin
- Introduced after the discovery of ribozymes
 - RNA molecules with catalytic properties
- According to this theory, the genetic system is the first to emerge, but with self-catalytic properties
 - Present-day ribozymes would be a sort of molecular fossiles of an ancient “RNA world”



Walter Gilbert



Emergence of the “DNA world”

- Present-day DNA-world would have emerged at a later stage because of its advantages:
 - DNA is more stable
 - The lack of an oxygen atom in the sugar (deoxyribose instead of ribose) makes DNA less reactive than RNA
 - The double DNA strand is more stable than the single RNA strand
 - DNA is better preserved because it is only involved in transcription (not in translation)
 - In eukaryotes DNA is preserved inside the nucleus of the cell
 - The DNA world has an extremely greater flexibility
 - Due to the introduction of proteins specialized in a large variety of metabolic functions

Life as a kinetic state of matter

Addy Pross


Example of the kinetic power of self replication

- Comparison between normal and self-catalytic reactions
 - start with 1 molecule of catalyst X
 - assume reaction rate $1\mu\text{s}$ in both cases
- Time required to build up a mole of products (6×10^{23})
 - Normal case: 20 billion years
 - Self-catalytic case: $79 \mu\text{s}$
- The kinetic control of chemical reactions could be the key for understanding the origin of life (in chemistry, the term “kinetics” is related to the rate of chemical reactions)
 - see literature by Addy Pross


A, B : reactants

X : catalyst

Normal case



Self-catalytic case



Replication and molecular evolution

Imperfect replication and chemical selection are supposed to be the key ingredients of some form of molecular evolution that has supposedly led to the molecular machinery that we see today

Molecular replication and chemical selection is a conceptual extension back in time, down to the prebiotic stages, of the concept of Darwinian evolution of living species (reproduction and natural selection)

The capability of replication is probably the key function for the initial selection of biomolecules

Replication and molecular evolution

Evolution driven by natural selection,
whether of species or molecules,
is not teleological

In other words, it is not the result of plan

Evolution driven by natural selection works *a posteriori*,
favouring the most suitable variations
for a given function that already exists

Compartments

In order to develop protocells, the early products of the RNA-world must have been enclosed in compartments

Compartmentalization is required to prevent the dispersion of genetic information and to concentrate the action of cooperative biochemical processes in an enclosed space

Membranes delimit a set of structures and reactions that can be transmitted as a specific heritage, paving the road for the onset of Darwinian evolution

Early membranes

In present-day life the compartments are provided by the phospholipid bilayers of the cell membranes

Phospholipids are the result of an evolutionary process, and their synthesis requires enzymatically catalyzed reactions not available for the first protocells

Early membranes could have been constituted by simple fatty acids

Simple fatty acids can be spontaneously generated in prebiotic chemistry, as demonstrated, for example, by their presence in the Murchison meteorite





Figure credits: Di Mauro & Saladino

Protocellular vesicles

Laboratory experiments demonstrate that simple amphiphilic molecules, resulting from prebiotic processes, can give rise to vesicle structures

Variations of the *concentration* of amphiphilic molecules and of *ambient conditions* drive the formation and destruction of vesicle structures that can grow and duplicate


Jack Szostak demonstrated that protocellular vesicles better replicate if they contain RNA and, at the same time, RNA better replicates if it is enclosed in lipidic vesicles



Casting light on the first living cells

- “Top-down” approach
 - From the study of present-day living organisms, we try to characterize the properties of the first terrestrial organisms proceeding backwards in evolution
- The top-down approach is based on the comparison of genetic sequences of different species
 - The techniques of molecular biology allow us to classify species on the basis of their genetic sequences, rather than on their morphology or phenotype
 - Phenotype: composite of observable traits and behaviour of organisms
 - The results are visualized in a “phylogenetic tree”, where the distances between different species are proportional to the differences found in the genetic sequences

The phylogenetic tree of life




The classification based on genetic sequences has revolutionized our understanding of unicellular organisms

Through genetic classification we have been able to distinguish three domains of unicellular organisms:

Archaea, Eubacteria and Eukaryotes

Archaea were previously classified as bacteria




A recent view of the phylogenetic tree of life (Hug et al. 2016)

Bacteria are predominant over archaea and eukaryotes

Archaea are closer to Eukaryotes than Bacteria

The Last Universal Common Ancestor




LUCA = Last Universal Common Ancestor of present-day living organisms, also called Cenancestor

Close to the "root" of the tree, we find thermophilic Archaea and Bacteria

The gap between the RNA world and the LUCA


- The root of the phylogenetic tree is not representative of the oldest living cell
 - Other forms of life, extinct in the course of the evolution, must have preceded the LUCA
 - The earliest form of life is sometimes called the “progenote”
 - The early life could have been a collection of somewhat different cells, rather than a single type of cell
 - Detailed analysis suggests that *early life was mesophilic, rather than thermophilic*



Horizontal gene transfer

(also called Lateral Gene Transfer)

- Bacteria can exchange genetic material not only during their reproduction (“vertical gene transfer”, VGT) but also via direct exchange from one cell to another (“horizontal gene transfer”, HGT)
- The existence of HGT complicates the reconstruction of the phylogenetic tree, which is based on the VGT scenario
- HGT must have played an essential role in the early stages of life, providing a simple mechanism to exchange genetic material before more complex mechanisms of “vertical” transmission were set in place




Filling the gap from the RNA world to protocells

[FROM RNA WORLD TO BACTERIA]

Journey to the Modern Cell

After life got started, competition among life-forms fueled the drive toward ever more complex organisms. We may never know the exact details of early evolution, but here is a plausible sequence of some of the major events that led from the first protocell to DNA-based cells such as bacteria.




1 EVOLUTION STARTS ▲

The first protocell is just a sac of water and RNA and requires an external stimulus (such as cycles of heat and cold) to reproduce. But it will soon acquire new traits.

2 RNA CATALYSTS ▼

Ribozymes—folded RNA molecules analogous to protein-based enzymes—arise and take on such jobs as speeding up reproduction and strengthening the protocell's membrane. Consequently, protocells begin to reproduce on their own.




3 METABOLISM BEGINS ▲

Other ribozymes catalyze metabolism—chains of chemical reactions that enable protocells to tap into nutrients from the environment.


Ricardo & Szostak (2009)

Filling the gas from protocells to prokaryotes



Ricardo & Szostak (2009)

Possible scenarios for the origin of life



Autotrophic:


the first organisms extract energy and synthesise organic material from the abiotic world

Heterotrophic:

the first organisms harvest organic material and energy from prebiotic molecules that are already present in the environment


The heterotrophic scenario

The molecular ingredients could have been synthesized on the primitive Earth and/or could have been delivered on Earth from space



This hypothesis does not require a specific environmental niche:
different types of subaerial sites can in principle be involved

The heterotrophic scenario



- **Anoxic geothermal fields**

- Geothermal fields are conducive to condensation reactions and enable the involvement of solar light as an energy source and as a selector factor of stable nucleotides
- Supported by geochemical data and phylogenomic analysis (Mulkidjanian et al. 2012)
- Geothermal vapour is enriched in phosphorus compounds that could be essential for the emergence of the first RNA-like oligomers

Comparison of heterotrophic and autotrophic scenarios

Planetary environments suitable for abiogenesis

Scenario of abiogenesis	Variety of sites or conditions	Transfer of prebiotic products	Variations in a given site	Exogenous delivery	Origin of chirality
Subaerial sites (heterotrophic scenario)	Ponds, lakes, beaches, dry areas, volcanoes, geysers, hot springs	PLAUSIBLE (e.g., hydrological system)	Irradiation (day-night cycle), concentration (dry-wet cycles in ponds/lakes, tidal cycles on beaches)	POSSIBLE	On Earth's surface or from meteorites
Bottom of the oceans (autotrophic scenario)	VERY LIMITED (inside vent system)	VERY LIMITED (inside vent system)	NO (steady conditions)	NO (extremely diluted)	NO