

Astrobiology

Lecture 13

The inner Solar System

Trieste University, Academic Year 2021-2022

Giovanni Vladilo (INAF-OATs)

Search for habitable planets and life in the Solar System

Most studies of the Solar System are based on results obtained from interplanetary space missions

These investigations are complemented by laboratory experiments and studies of extremophiles simulating planetary space conditions

Venus

Surface conditions

$$T_s = 735 \text{ K}$$


$$p_s = 92 \times 10^5 \text{ Pa}$$

Atmospheric composition dominated by CO_2 , without O_2

Absence of tectonics

Surface conditions do not satisfy the liquid water criterion: currently non habitable

Venus has probably undergone a “runaway greenhouse effect” in the early stages of its history



Venus


Origin of the CO₂-rich, dense atmosphere

Possible scenarios:

- As a result of a runaway greenhouse instability, water would be photodissociated and hydrogen lost, leaving free oxygen that would combine with carbon (the absence of hydrogen inhibits the chemical pathways that lead to organic chemistry, e.g. CH₄)
- The atmosphere was CO₂-rich and dense since the beginning; in this scenario, the loss of water through runaway greenhouse instability prevents the formation of oceans and the possibility of tectonics; without oceans and without tectonics it is not possible to remove CO₂ from the atmosphere


Searching for “habitable” layers in the atmosphere of Venus

- At a height of ~50-60 km, p and T lie within the liquid water limits, being roughly similar to the values found at the surface of the Earth
- Based on the liquid criterion alone, life could in principle exist



Claims of detection of phosphine in Venus atmosphere

- Recent work have claimed the detection of phosphine (PH₃) in the atmosphere of Venus (Greaves et al. 2020)



- No known abiotic process would be able to generate phosphine in terrestrial-type planets
- Claims that phosphine could be generated by biological processes have been advanced
- However, the detection of phosphine has been strongly criticized and is currently not confirmed
- Also from the theoretical point of view, the existence of life in confined layers of the Venusian atmosphere is quite challenging
- Due to strong winds and convective motions no type of material would be able to remain suspended in the “habitable” layer

Mars

Surface habitability

At present time the Mars surface is not habitable

The surface is slightly below the triple point of water (611 Pa)

$$T_s = 210 \text{ K}$$



$$P_s \sim 600 \text{ Pa} (\sim 6 \text{ mbar})$$

The atmospheric composition is dominated by CO_2

Tectonics is absent

At some proper depth below the surface we expect conditions suitable for liquid water due to pressure and temperature gradients

Salinity would help to decrease the freezing point




Search for water in Mars

The search for water has been one the main astrobiological goals of Mars studies

Evidence of water in present-day Mars

Traces of recent erosion at the border of craters

Interpreted as transient outflows of water in liquid phase (“gullies”)



Mars Reconnaissance Orbiter, NASA (2006)

Search for water in Mars

Evidence of water in present-day Mars

The bulk of the polar caps is constituted by CO_2 ice, but the North polar cap must also contain H_2O

this would explain why such polar cap is able to persist, to some extent, during the Mars summer, when CO_2 sublimates into the atmosphere



Search for water in Mars


Evidence of water in present-day Mars

Space probes are collecting evidence of underground water ice

The distribution of hydrogen below the ground, inferred from the data collected from the probe “Mars Odyssey”, suggests the existence of a layer of water ice at a depth of about one meter


Blue areas: maximum concentration

Mars Odyssey, NASA (2001)



Radar evidence of subglacial liquid water on Mars

South Polar Layered Deposits



Reflected signal interpreted as evidence of a stable body of liquid water at ~1.5 km below the surface

Orosei et al. (2018)

Mars habitability in the past

The evidence for water in Mars is compelling, but at present time water seems to be largely confined in ice phase below the surface, with sporadic signatures of outflows and sublimation

However, the situation must have been very different in the past of Mars

Several independent evidences suggests that Mars was habitable in the past

These evidences are particularly important in astrobiology, because of the possibility that life might have emerged on Mars

Among the different evidences of past habitability we mention:

Statistics of impact craters

Geomorphological evidence

Martian meteorites collected on Earth

Mars habitability in the past

Evidence for the presence of a thick atmosphere in the past

Statistics of the diameters of impact craters

Deficit of small size ancient craters with respect to recent craters

The presence of a past atmosphere may have caused this deficit by means of:

Fusion of small size meteoroids due to friction during the crossing of the atmosphere

Erosion of the shallowest craters by means of atmospheric weathering

A thick atmosphere would have dramatically enhanced the habitability of Mars:

increasing the pressure above the triple point of water


increasing the greenhouse effect and surface temperature

protecting the planetary surface from ionizing radiation

Mars: evidence for past geological activity

Tectonic fractures, ridge-like shapes

These features suggest that episodes of fluid alteration along the fractures must have taken place in past geological times



Mars Reconnaissance Orbiter, NASA (2006)



Mars habitability in the past

Evidence for the presence of liquid water in the past

Geomorphological evidence

Network of valleys similar to those excavated by terrestrial rivers

Mars experienced at least short periods of clement conditions towards the end of the Noachian Era (~4.1 to 3.7 Gya) that supported a hydrological cycle



Mars habitability in the past

Evidence for the presence of liquid water in Mars meteorites

Some meteorites recovered in Allan Hills (Antarctica), are of martian origin and suggest that liquid water was present on Mars in its early geological eras

SNC meteorite ALH 84001 (found in 1984)

Radiodating: 4.1 Gya (Lapen et al. 2010)




Mars habitability in the past

- In light of the “faint young sun paradox”, the requirement for a primordial CO₂-rich atmosphere is more compelling for Mars than Earth
- An intense, early volcanic activity may have generated large amounts of atmospheric CO₂

The greenhouse effect would have provided a temperature sufficiently high for water being in the liquid phase

However, the atmospheric pressure should have been much higher than today (several bars of CO₂)



Atmospheric loss in early Mars

The large amount of atmospheric CO₂ required to solve the faint young paradox for Mars is not confirmed by the geochemical evidence of Mars surface


It is possible that Mars atmosphere was quickly dissolved by intense solar wind at early stages of Mars evolution

The Mars Atmosphere and Volatile Evolution (MAVEN) spacecraft has been orbiting Mars since 2014 to address this problem

This mission aims at characterizing the upper atmosphere and ionospheric structure and composition, the interactions of the sun and the solar wind, and the processes driving loss of atmospheric gas to space (Jakosky et al. 2015)

Results from the MAVEN mission indicate that major episodes of solar activity are extremely efficient at driving loss of Mars atmosphere

Searches for life on Mars



Viking experiments (1976) searched for traces of biological activity from the analysis of samples collected in a few martian landing sites

Several experiments were carried out and analysed *in situ*, searching for evidence of biochemical processes

A signal from the Labeled Release (LR) experiments was consistent with the presence of biochemical activity, but this signal is generally believed to be a “false positive” because it was not confirmed by the Gas Chromatograph - Mass Spectrometer (GCMS) experiment

Lessons from the Viking experiment

The ambiguous results of the Viking experiments teach us how difficult is to reveal the presence of life, even when we can analyse samples

Analysis of Martian samples in Earth laboratories would be much more accurate than the analysis performed in situ, but bringing the sample back to Earth would increase dramatically the cost of the mission

The negative (or ambiguous) results of the Viking experiments do not exclude that life might exist in other locations on Mars

Active biological processes might take place in underground layers at a proper depth, where the temperature and pressure gradients would allow liquid water to be present

If so, we would expected to find some biosignatures, such as traces of surface gas of biological origin

Search for atmospheric biosignatures in Mars

The atmospheric composition is dominated by CO₂

Given the low value of partial pressure, the greenhouse effect driven by CO₂ is insufficient to make Mars habitable at present time

O₂, the most classic atmospheric biomarker on Earth, is absent in the atmosphere of Mars

Detections of traces of CH₄ in the Mars atmosphere have been reported several times

Methane emissions seem to have a local and seasonal character

If confirmed, outgassing of CH₄ would suggest the presence of underground chemical activity

The activity could be geochemical or even biochemical, based on the biogenic production of methane on Earth

The search for atmospheric methane in Mars


Claims of detection:

- remote observations of a gas plume (Mumma et al. 2009)
- occasional detections by NASA's Curiosity rover
- ESA's Mars Express spacecraft (Giuranna et al. 2019)

Unsuccessful search:

Trace Gas Orbiter (European-Russian)

Methane undetected at a height of 5 km from the surface





Criticism

CH₄ could be real, but may result from contamination of terrestrial rovers
(Kevin Zahnle)


Searches for life in the past history of Mars

The analysis of the meteorite ALH 84001 revealed microstructures with morphology suggestive of a biological origin

Carbonate globules with an age of 3.9 Ga have been found in its interior
However, the sizes of those structures, between 20 and 100 nm, are too small with respect the smallest sizes of the living cells that we know



Mars contamination



Different planetary protection cleanliness levels for different parts of a spacecraft do not necessarily prevent soil contamination because these cleaning strategies evolved without consideration of the effects of the descent engine plumes.